




Survey of the Status of Important Fauna Species in the Kyparissiakos Lease Area

Final Report



November 2024

[This page intentionally left blank]

Contents

Contents	i
1 Introduction	5
2 Description of the Project Area	6
2.1 Oceanographic characteristics of the Project Area	8
2.2 General information of the main cetacean, seabird and sea turtle species in the Project Area	10
2.2.1 Cetaceans	10
2.2.2 Seals	16
2.2.3 Sea turtles	16
2.2.4 Seabirds	20
3 Methodology	24
3.1 Pelagic surveys	24
3.1.1 Boat surveys	24
3.1.2 Aerial surveys	28
3.2 Coastal surveys	30
3.2.1 Coastal surveys for the Scopoli's Shearwater	30
3.2.2 Coastal surveys for the Mediterranean Shag	31
3.2.3 Coastal surveys for the Mediterranean Monk Seal	31
Cave monitoring with the use of Automated Infrared Cameras	31
3.3 Sea turtle telemetry	32
3.4 Telemetry of Scopoli's Shearwater	33
4 Results	35
4.1 Pelagic surveys	35
4.1.1 Boat surveys	35
4.1.2 Aerial surveys	37
4.2 Coastal surveys	50
4.2.1 Coastal surveys for the Scopoli's shearwater	50
4.2.2 Coastal surveys for the Mediterranean Shag	50
4.2.3 Coastal surveys for the Mediterranean Monk Seal	50
4.3 Sea turtle telemetry	52
4.4 Telemetry of Scopoli's Shearwater	54
5 Conclusions	56
Bibliography	57

Annex I: Report on the coastal surveys for the Mediterranean monk seal	64
Annex II: Report on Loggerhead turtles tracking in Kyparissia	64

Project Team

Name	Company	Expertise	Role
Tasos Dimalexis	NCC Ltd.	PhD Biologist, Ornithologist	Coordinator, field & drone surveys
Thanos Kastritis	NCC Ltd.	PhD Oceanographer, Ornithologist	Field surveys, Reporting
Manolia Sergi	NCC Ltd.	Biologist, Certified Marine Mammal Observer, Certified Passive Acoustic Monitoring Operator	Marine Mammal Observer, PAM Operator
Jonathan Gordon	NCC Ltd.	PhD Zoologist, University of St Andrews, Sea Mammal Research Unit, Research Fellow	Cetaceans Expert, field surveys
Niki Koutouzi	NCC Ltd.	MSc Oceanographer	Marine Mammal Observer, PAM Operator
Eirini Antoniadi	NCC Ltd.	MSc Biologist	Marine Mammal Observer, PAM Operator
Giorgos Kouthouridis	NCC Ltd.	Field Ornithologist	Field surveys
Alan Rees	NCC Ltd	PhD Biologist/Sea Turtle Expert	Sea Turtle Satellite Tagging Leader
Odysseas Stamatakis	NCC Ltd	Electronics Technician	Drone operator
Vassilis Goritsas	NCC Ltd.	Environmental MSc	GIS & Data management
Panagiotis Dendrinou	MOM/BIOTOPIA	PhD Marine Biologist	Coordinator, Marine and Aerial Surveys Expert
Kimón Koemtzipoulos	MOM	MSc Marine Biologist, Certified Marine Mammal Observer	Marine Mammal Expert/ Certified MMO, Field research & Data compilation
Odysseus Paxinos	MOM	MSc Oceanographer	Sea Turtle Satellite Tagging, Monk Seal Field research & Data compilation
Panagiotis Mpatzios	MOM	System Engineer/Field Technician	Field Research and technical coordination
Styliani Adamantopoulou	MOM/BIOTOPIA	Biologist/Marine Mammal Expert	Data analysis, Reporting
Antonis Voutsinos	NCC Ltd.		Boat skipper
Kostas Papakonstantinou	NCC Ltd.		Boat skipper

Abbreviations and scientific names

<i>Calonectris diomedea</i>	Scopoli's Shearwater
<i>Caretta caretta</i>	Loggerhead Turtle
<i>Chelonia mydas</i>	Green Turtle
<i>Delphinus delphis</i>	Short-beaked Common Dolphin
<i>Grampus griseus</i>	Risso's Dolphin
<i>Hydrobates pelagicus</i>	European Storm-Petrel
ESAS	European Seabirds At Sea (survey method)
<i>Larus audouinii</i>	Audouin's Gull
<i>Larus michahellis</i>	Yellow-legged Gull
<i>Monachus monachus</i>	Mediterranean Monk Seal
n.m.	nautical mile
<i>Phalacrocorax aristotelis desmarestii</i>	Mediterranean Shag
<i>Physeter macrocephalus</i>	Sperm Whale
<i>Puffinus yelkouan</i>	Yelkouan Shearwater
<i>Stenella coeruleoalba</i>	Striped Dolphin
SAC	Special Area of Conservation (Natura 2000 network)
SPA	Special Protection Area (Natura 2000 network)
SDF	Standard Data Form (Natura 2000 datasheet)
<i>Tursiops truncatus</i>	Common Bottlenose Dolphin
WP	Work Package
<i>Ziphius cavirostris</i>	Cuvier's Beaked Whale

1 Introduction

In the context of Environmental Monitoring and Recording of Critical Environmental Indicators of Biodiversity, such as marine mammals (cetaceans and monk seals), sea turtles and seabirds, the Hellenic Petroleum Exploration & Production of Hydrocarbons Kyparissiakos Gulf Single Member S.A. company has assigned to Nature Conservation Consultants (NCC) Ltd a contract for conducting the present Project, namely the “Survey of the Status of Important Fauna Species in the Kyparissiakos Lease area” (Block 10).

The Project consists of 3 work packages (WP):

- I. **Pelagic Surveys for marine mammals, seabirds, sea turtles, nearshore and in the open sea**, using an open water RIB vessel, a sailing boat and a single engine aircraft.
- II. **Coastal surveys for monk seals and Mediterranean shag breeding sites in the coastal zones of the adjacent Natura 2000 sites**, using inflatable RIB boats.
- III. **Telemetry for seabirds and marine turtles** using drone videography and stationary thermal cameras, as well as satellite transmitters.

The present document consists of the **Final Report** of the **Work Packages WP I-III**. It presents the field surveys carried out during 2024 (April to October 2024) and the results in each Work Package of the project “Survey of the Status of Important Fauna Species in the Kyparissiakos Lease area”.

The present project is the 2024 continuation of the ongoing project “Survey of the Status of Important Fauna Species in the Kyparissiakos Lease area”, implemented in the period 2020-2023.

2 Description of the Project Area

The **Project Area** is located in the Ionian Sea, southeast of Zakynthos Island and west of Peloponnese, approximately from the latitude town Zacharo in the north and town Methoni in the south. It extends between latitudes of $36^{\circ}50'N$ in the south and $37^{\circ}30'N$ in the north and between longitudes of $20^{\circ}55'E$ in the west and $21^{\circ}30'E$ in the east. Its total surface area is $3,422.5 \text{ km}^2$ (Figure 2-1).




Figure 2-1. Project Area

The **Wider Project Area** envelops the project area and extends further north and east to additionally include the southwestern, south-eastern and eastern coast of Zakynthos, and the western coast of Peloponnese south of Kyllini, together with their neighbouring islets (Figure 2-2). The **Pelagic Survey Area**, where pelagic surveys are carried out includes the primarily the Project Area together with neighbouring waters to the east.




Figure 2-2. Wider Project Area

2.1 Oceanographic characteristics of the Project Area

The sea depth within the Project Area exceeds 500m and reaches more than 3,500m at its southwestern corner. The only exception are the Strofades islets in the west, which are surrounded by a narrow belt of coastal waters (Figure 2-3).




Figure 2-3. Bathymetry in the Project Area

In northern, north-eastern and eastern part of the Wider Project Area the slope of the sea floor descends gradually, without abrupt breaks towards southwest, however the southern and the western part exhibit numerous abrupt descends of the sea floor, accompanied by step slopes of the sea floor reaching up to 53° (Figure 2-4).




Figure 2-4. Slope of the sea floor in the Wider Project Area

2.2 General information of the main cetacean, seabird and sea turtle species in the Project Area

2.2.1 Cetaceans

Hellenic seas host an unexpectedly high diversity of cetaceans with eight (8) species that are resident in the area, seven (7) of which belong to the Odontoceti suborder: Sperm Whale (*Physeter macrocephalus*), Cuvier's Beaked Whale (*Ziphius cavirostris*), Risso's Dolphin (*Grampus griseus*), Bottlenose Dolphin (*Tursiops truncatus*), Striped Dolphin (*Stenella coeruleoalba*), Short-beaked Common Dolphin (*Delphinus delphis*) and Harbour Porpoise (*Phocoena phocoena*) along with one representative of the Mysticeti suborder: Fin Whale (*Balaenoptera physalus*). The Harbour Porpoise is restricted to the Thracian Sea and North Aegean Sea, while the others are present one or more seas in Greece (Frantzis et al. 2003).

It is important to note that due to the semi-enclosed nature of the Mediterranean basin, in combination with its very particular oceanographic features and oligotrophic waters especially moving towards the east of the basin, cetacean species populations of the Mediterranean (which occur elsewhere in the world also) are treated separately by the IUCN, when it comes to the designation of their threat status and population trends. In the majority of cases, the Mediterranean subpopulation of cetacean species have at least one level higher in their designated threat status than the global population for the same species or are classified as Data Deficient.

The Wider Project Area is located along the Hellenic Trench, which is one of the most important areas for cetaceans in Greece. With the exception of the Harbour Porpoise (found only locally in the north-eastern Aegean) and the Fin Whale (observed mainly in the Ionian Sea, along the Hellenic Trench, north of Kefallonia), the remaining 6 commonly occurring species of cetaceans inhabiting Greek waters have been sighted or recorded as stranding in the Wider Project Area.

Table 2-1. General types of habitats, bathymetric characteristics and distance from coast of recorded presence in Greek seas of common cetacean species that are present in the Wider Project Area (from Frantzis 2009).

Species	Common name	Habitat		
		Type	Depth	Distance from coast
<i>Physeter macrocephalus</i>	Sperm whale	Slope, secondarily pelagic	1235 m (510-2933 m)	8.1 km (1.6-25.2 km)
<i>Ziphius cavirostris</i>	Cuvier's beaked whale	Slope, probably pelagic as well	1066 m (491-2279 m)	8.6 km (2.1-26.5 km)
<i>Grampus griseus</i>	Risso's dolphin	Slope, probably over its shallower part	737 m (165-1717 m)	8.2 km (0.3-28.3 km)
<i>Tursiops truncatus</i>	Common bottlenose dolphin	Typically, coastal, also over shallow waters "offshore"	121 m (1-1504 m)	3.0 km (0.0-26.0 km)
<i>Stenella coeruleoalba</i>	Striped dolphin	Typically, pelagic and slope	1024 m (75-2920 m)	8.7 km (0.6-37.1 km)
<i>Delphinus delphis</i>	Short-beaked Common dolphin	Coastal and shallow, ("pelagic" and deep only in the Gulf of Corinth)	86 m (11-274 m) Gulf of Corinth: 713 m (275-935)	8.7 km (0.6-37.1 km)

The Wider Project Area includes, coastal areas, continental shelf and slope, as well as pelagic areas. For the purpose of the present study and based on the types of marine habitats typically used by the species present in the Wider Project Area, the focus of pelagic surveys is primarily on the species with regular presence in the Wider Project Area, namely the **Sperm Whale (*Physeter macrocephalus*)**, **Cuvier's Beaked Whale (*Ziphius cavirostris*)**, **Striped Dolphin (*Stenella coeruleoalba*)** and **Risso's dolphin (*Grampus griseus*)** in the pelagic and continental slope areas, and **Short-Beaked Common Dolphin (*Delphinus delphis*)** and **Bottlenose Dolphin (*Tursiops truncatus*)** in coastal areas. Accounts on the biology, ecology, as well as conservation and threat status of the cetacean species of interest are provided below. It should be noted that large data gaps are still present regarding the distribution and abundance of cetaceans in the eastern Mediterranean (Mannocci et al. 2018).

2.2.1.1 Sperm Whale (*Physeter macrocephalus*)



Figure 2-5. Sperm Whale (*Physeter macrocephalus*) (© Massimo Demma/ICRAM)

The second largest cetacean found in Greece and the largest Odontocetus found globally is the Sperm Whale (*Physeter macrocephalus*). The Sperm Whale prefers deep water habitats particularly deep continental slope water where they hunt their preferred prey, large mesopelagic cephalopods (Frantzis 2009, Notarbartolo di Sciara et al. 2012).

The Hellenic Trench is considered to be the species core habitat for the eastern Mediterranean sub-population (Frantzis et al. 2014). The total species population size in the Greek Seas is estimated at 180 – 280 individuals (2013-18 Habitats Directive Article 17 Reporting at <https://nature-art17.eionet.europa.eu/article17/>), the population size in the Hellenic Trench 200 – 250 individuals (Frantzis et al. 2014) and the estimated population size in the Ionian Sea, including international and Italian waters 62 individuals (95% CI: 24-165 individuals, in Lewis et al. 2003), however this is likely to be an underestimation (Frantzis 2009).

2.2.1.2 Cuvier's Beaked Whale (*Ziphius cavirostris*)




Figure 2-6. Cuvier's Beaked Whale (*Ziphius cavirostris*) (©Artescienza-M.Würtz)

Cuvier's Beaked Whale, a medium sized odontocetus, shares the same habitat and distribution as that described for the Sperm Whale, namely the continental slope. Almost all past species sightings occurred above depths of 500-1,500m (Frantzis et al. 2003). It is the only beaked whale common in the Mediterranean Sea. In Greece, the majority of past sightings are associated with the Hellenic Trench, from eastern Rodos Island to northwest Corfu Island (Frantzis et al. 2003, Frantzis 2009) with the highest number of sightings south of Crete and west of Lefkada (Frantzis et al. 2003, Podestà et al. 2016). Along the Hellenic Trench the species feeds almost exclusively on mesopelagic and bathypelagic cephalopods (Frantzis

2009). Several sightings and numerous strandings have been recorded in the Wider Project Area (based on Frantzis 2009).

The Hellenic Trench is one of the species high-density areas in the Mediterranean. The total species population size in the Greek Seas as well as in the Wider Project Area is unknown (2013-18 Habitats Directive Article 17 Reporting at <https://nature-art17.eionet.europa.eu/article17/>). It is worth noting that Greek seas are considered to host quite a significant portion of the Mediterranean population (Frantzis 2009).

2.2.1.3 Risso's Dolphin (*Grampus griseus*)



Figure 2-7. Risso's dolphin (*Grampus griseus*) (© Massimo Demma)

Risso's dolphin is the largest dolphin that commonly occurs in the Greek Seas. The sightings and strandings records indicate that the species is present in all parts of the Greek Seas, however the only known area where the species is predictably present is the Myrtoon Sea extending south to the north-western Crete. The species is present in the Ionian Sea, as confirmed by strandings which have been recorded from north Corfu Island to south Peloponnese. No sighting records have been made in the Ionian Sea which indicates that either the species is present in low numbers or it is present outside warm period when past surveys have been made. The strandings in the Ionian Sea have been recorded from the end of September until late April. The species is present primarily along the continental slope, preferably deep water and shelf break where the slope is the steepest, but also close to the coast, particularly when the shelf is narrow (Frantzis 2009). The species feeds mainly with squid and occasionally with fish.

The total species population size in the Greek Seas is estimated to be 100 – 600 individuals (2013-18 Habitats Directive Article 17 Reporting at <https://nature-art17.eionet.europa.eu/article17/>). The population size in the in the Wider Project Area is unknown.

2.2.1.4 Bottlenose dolphin (*Tursiops truncatus*)




Figure 2-8. Common bottlenose dolphin (*Tursiops truncatus*) (© Artescienza-M. Würtz)

The bottlenose dolphin is the most common species of dolphin found in coastal shallow waters of the Mediterranean (Frantzis 2009). It is homogeneously distributed across all Greek Seas as it has been sighted in most coastal areas, straights and gulfs. (Frantzis 2009). The Bottlenose Dolphin in Greece, similar to Short-beaked Common Dolphin prefers the continental shelf usually staying within a depth of up to 200m (Frantzis 2009). It is known to consume a variety of prey items being quite adaptive.

The total species population size in the Greek Seas is estimated to be 3,800 – 9,000 individuals (2013-18 Habitats Directive Article 17 Reporting at <https://nature-art17.eionet.europa.eu/article17/>). The population size in the Wider Project Area is unknown.

2.2.1.5 Striped dolphin (*Stenella coeruleoalba*)




Figure 2-9. Striped dolphin (*Stenella coeruleoalba*) (© Massimo Demma/ICRAM)

The Striped Dolphin, a small delphinid, has a year-round presence in Greek waters. It is the most abundant dolphin species in Greece and the Mediterranean overall (Frantzis 2009). Its distribution in Greece is widespread and it occurs in all deep (>500m), pelagic waters and the continental slope but it can also inhabit intermediate depths of 200-500m (Frantzis 2009). The Striped Dolphin is frequently sighted along the length of the Hellenic Trench. The species diet includes mainly cephalopods, as well as fish and crustaceans.

The total species population size in the Greeks Seas is estimated to be 20,000 – 80,000 individuals (2013-18 Habitats Directive Article 17 Reporting at <https://nature-art17.eionet.europa.eu/article17/>). The population size in the in the Wider Project Area is unknown.

2.2.1.6 Short-beaked common dolphin (*Delphinus delphis*)




Figure 2-10. Short-beaked common dolphin (Delphinus delphis) (© Artescienza-M. Würtz)

The Short Beaked Common Dolphin (or simply Common Dolphin) is a small delphinid with a year-round presence in Greek waters. Its distribution in Greece is patchy and their presence seems to be mostly limited to the central and northern Greek Seas (Frantzis 2009). In general, it prefers shallow (<200m) and coastal waters, with exception of Gulf of Corinth where it exhibits preference to pelagic habitats (Frantzis 2009). It exhibits flexible feeding habits. The distribution of the Common Dolphin in the Ionian Sea the is limited to shallow waters between north Lefkada, Kefallonia and south Zakynthos and the mainland. In the Inner Ionian Sea, the main prey includes shoaling fish e.g., anchovies and sardines.

The total species population size in the Greeks Seas is estimated to be 750 – 4,200 individuals (2013-18 Habitats Directive Article 17 Reporting at <https://nature-art17.eionet.europa.eu/article17/>).

The population of Common Dolphins of the Inner Ionian Sea has been the focus of regular surveys for years and has been well documented (Bearzi et al. 2008B). The local population counted 150 individuals until the mid-90s and their range seemed to cover the entire Inner Ionian. Since then, the population has declined dramatically with only an estimated 15 individuals encountered over the past years mostly sighted in southern Lefkada (Bearzi et al. 2008B).

2.2.2 Seals

2.2.2.1 Mediterranean Monk Seal



© G. Gousis/MOM

Figure 2-11. Striped dolphin (*Stenella coeruleoalba*) (© Massimo Demma/ICRAM)

The Mediterranean Monk Seal is the only pinniped (seal) living in the Mediterranean region, the rarest extant member of the Phocidae family and one of the rarest marine mammals in the world.

Mediterranean monk seals were once widely and continuously distributed in the Mediterranean and Black Seas, and in the North Atlantic waters from Morocco to Cap Blanc, including the Canary, Madeira and the Azores Islands. A few individuals have been recorded in Senegal, the Gambia and the Cape Verde Islands in the southern end, as well as in Portugal and Atlantic France in the northern end of the species' distribution. Today the distribution of the Mediterranean is highly fragmented and consists of three to four isolated subpopulations (Karamanlidis et al. 2016). In the Mediterranean Sea, the stronghold of the species has been on islands in the Ionian and Aegean Seas, and along the coasts of Greece and western and southern Turkey ((Güçlüsoy, Kiraç, Veryeri, & Savaş 2004, Gücü, Gücü, & Orek 2004, Anonymous, 2007). In the North Atlantic, two subpopulations exist: one at Cabo Blanco (also known as Cap Blanc) at the border of Mauritania and Western Sahara (González & Fernandez de Larrinoa 2012, Martínez-Jauregui et al. 2012), and one at the Archipelago of Madeira (Pires, Neves, & Karamanlidis, 2008). An unknown number of monk seals might still survive at the Mediterranean coasts of eastern Morocco (and perhaps Algeria) (Mo, Bazairi, Bayed, & Agnesi, 2011), but without on-going systematic conservation actions the fate of this subpopulation is unknown.

The total species population size in the Greece is estimated to be 300 – 400 individuals (2013-18 Habitats Directive Article 17 Reporting at <https://nature-art17.eionet.europa.eu/article17/>).

2.2.3 Sea turtles

There are three species of sea turtles that regularly occur in the Mediterranean: **Loggerhead Turtle (*Caretta caretta*)**, **Green Turtle (*Chelonia mydas*)** and **Leatherback sea turtle (*Dermochelys coriacea*)**. The sea turtles live almost exclusively in the marine environment with females returning to land for dig nests and lay eggs, while males almost never return to

land. The range of all three species extends along the Wider Project Area (Legakis & Maragou 2009, 2013-18 Habitats Directive Article 17 Reporting: species range), however only Loggerhead Turtle and Green Turtle have been recorded in the area (2013-18 Habitats Directive Article 17 Reporting: species distribution). Among these two the Loggerhead Turtle is the species of interest due to its regular presence and nesting in the Wider Project Area of the Kyparissia Bay, while the Green Turtle is regular but rare visitor in the area. The Leatherback sea turtle is only considered in Greece to be a visitor from the Atlantic (Casale & Margaritoulis 2010).

2.2.3.1 Loggerhead turtle (*Caretta caretta*)



Figure 2-12. Loggerhead Turtle (*Caretta caretta*)

The Loggerhead turtle is an oceanic turtle with a global distribution. It is a migratory species and may travel thousands of kilometres to forage and to return to its breeding sites. After hatching, logger-head turtles adopt an oceanic lifestyle in major current systems (Bolten and Witherington 2003). After 4-19 years spent in the oceanic zone, they move to neritic areas where they forage and mature over 10-39 years (Arens and Snover 2013). After attaining sexual maturity, they migrate between neritic foraging grounds and nesting areas. The Mediterranean, where the species is nesting in the eastern basin (Legakis & Maragou 2009), the breeding population of the loggerhead turtle is spread over tens of rookeries which are estimated to produce over 7,200 nests annually (Casale & Margaritoulis 2010) with the majority of nests being found in Greece. The country's two most important nesting beaches are located in the Wider Project Area, namely on Zakynthos (Laganas Bay) and on Peloponnese (Kyparissia Bay), which host 43% and 19% of all nests in Greece, respectively (Legakis & Maragou 2009). The average number of nests per season for the period 1984-2007 at Laganas Bay and at Kyparissia Bay are 1,244 nests/season (range: 833-2,018 nests/season) and 621 nests/season (range: 286-927 nests/season) (Casale & Margaritoulis 2010). Currently, Kyparissia Bay hosts the largest Loggerhead turtle nesting aggregation in the Mediterranean Sea (Rees et al. 2020).

In Greece and in the Central Mediterranean, the turtles after hatching disperse mainly in the Ionian, south-central Mediterranean and Adriatic Seas (Casale & Mariani 2014). Loggerhead turtles, especially juveniles, forage in almost all oceanic areas in the Mediterranean. Water circulation system has the greatest effect on their distribution (Casale et al. 2018). The neritic foraging areas (i.e., those located above continental shelf) are more frequently used by larger turtles, including adults (Casale et al. 2018, Figure 2-13). Loggerhead turtles generally

overwinter within or close to their foraging areas, however some may move from cold areas e.g., Adriatic Sea during winter (Casale et al. 2018).




Figure 2-13. Neritic foraging and wintering sites for loggerhead turtles (orange areas and arrows) and green turtle (green arrows) (adopted from Casale et al. 2018).

Migration corridors, are areas which are frequently used by migrating turtles, mainly for adult breeding migration and particularly for post-breeding migration from breeding areas to foraging grounds. Therefore, these migratory corridors are used at the end of the breeding season, in May and June by males, while in July and August, mostly by females (Casale et al. 2018). The main migration corridors are presented in Figure 2-14.




Figure 2-14. Main known migratory corridors for adult loggerhead turtles to and from breeding sites (stars). Light brown areas represent migratory funnels in the open sea while darker strips represent paths along the coasts, typically in shallow waters (adopted from Casale et al. 2018).

The movements of the Loggerhead turtles nesting in the Ionian Sea, particularly those from Zakynthos has been well studied by satellite or GPS telemetry (e.g., Zbinden et al. 2008, Schofield et al. 2010a-c, Schofield et al. 2013, Luschi & Casale 2014). The data from 75 tracked turtles breeding on Zakynthos showed after breeding the turtles migrate to neritic sites with waters shallower than 100m, with the majority of turtles migrate north to the Adriatic Sea and Amvrakikos Gulf (42%) or south-west to Libya and Tunisia (32%), while the remaining either

stay in the Ionian Sea or move to the eastern or western Mediterranean (Zbinden et al. 2008, Schofield et al. 2013). After leaving their foraging areas (in October – November) the tracked turtles move to their overwintering areas further south (Zbinden et al. 2008). The main foraging and overwintering areas are presented in the Map 11, below. The main foraging areas are located over the continental shelves and slopes (Ullmann & Stachowitsch 2015) in the Northern and Southern Adriatic Sea, Ionian Sea, the Strait of Sicily and the Tunisian shelf. A small proportion (~7%) were resident to Zakynthos. Significantly more males than females remain within 100km of Zakynthos (Schofield et al. 2013).

2.2.3.2 Green turtle (*Chelonia mydas*)




Figure 2-15. Green turtle (*Chelonia mydas*)

The green turtle (*Chelonia mydas*) is an migratory oceanic turtle with a global distribution. Their nesting sites in the Mediterranean are located mostly in Turkey, Cyprus and Syria (Figure 2-13) with an average of 1500 nests per year. No regular nesting areas are located in Greece. They use mostly marine areas in the Levantine basin, but also forage in Greece and Libya, as well as occasionally in the Adriatic Sea and the western Mediterranean basin (Figure 2-16). In Greece local concentration have been found in Lakonikos Bay, southern Peloponnese. Stranding data indicate that there is a more frequent presence of adult green turtles in southern Aegean (Casale & Margaritoulis 2010). The species has been recorded in the Wider Project Area.




Figure 2-16. Main known migration corridors for adult female green turtles during reproductive migrations from the breeding sites (stars) (adopted from Casale et al. 2018).

2.2.4 Seabirds

For the purpose of the present study, only those seabird species which are exclusively associated with the marine environment and the pelagic area, that have been recorded in the Ionian Sea in the past and their presence in the wider Project area has been either confirmed. These species include pelagic seabird species: **Scopoli's Shearwater (*Calonectris diomedea*)**, **Yelkouan Shearwater (*Puffinus yelkouan*)** and **European Storm-petrel (*Hydrobates pelagicus*)**, as well as, coastal seabird species which could be present in the pelagic areas due to shallow waters in the Project area or due to human activities, i.e. **Yellow-legged Gull (*Larus michahellis*)** and the **Mediterranean Shag (*Phalacrocorax aristotelis desmarestii*)**.

2.2.4.1 Scopoli's's Shearwater (*Calonectris diomedea*)




Figure 2-17. Scopoli's Shearwater (*Calonectris diomedea*) (© Paul Hirst)

Scopoli's Shearwater (*Calonectris diomedea*) breeds across Mediterranean with the majority of the population spending the non-breeding season in the Atlantic. In the past it was considered conspecific with the Cory's Shearwater (*Calonectris borealis*) which breeds in the Atlantic. In Greece the species breeding in the Aegean and Ionian Sea with the largest known colony being located at Strofades Islets (within the Project Area), south of the Zakynthos Island in the Ionian Sea, with an estimated breeding population of 5,550 pairs (Karris et al. 2017). Other large colonies occur mainly in the southern, central and eastern Aegean Sea although breeding has also been confirmed in the northern Aegean Sea (Fric et al. 2012). The only other known breeding area in the Ionian Sea is at Diapontia islands at Kerkyra with much smaller breeding population of 60-100 pairs (Fric et al. 2012).

2.2.4.2 *Yelkouan Shearwater (Puffinus yelkouan)*




Figure 2-18. *Yelkouan Shearwater (Puffinus yelkouan)* (© Paul Hirst)

Yelkouan Shearwater is an endemic species to the Mediterranean and the Black Sea. The known species colonies in Greece are located in the Aegean Sea, while no colonies have been found so far in the Ionian Sea. The main known colonies are located the North, East and Central Aegean Sea (Fric et al. 2012), with the largest being on Gyaros island in the Northern Cyclades (Fric & Portolou 2016). During the non-breeding season Yelkouan Shearwaters disperse widely within the Mediterranean Sea (mainly Adriatic and Aegean Seas) and the Black Sea. Additionally, 4,000-6,000 individuals are estimated to overwinter in the Aegean Sea. The main foraging areas of the Yelkouan Shearwaters are rich coastal and pelagic fishing grounds in the North, Central and East Aegean Sea, while the species is less common in the South Aegean and Ionian Seas (Fric et la. 2012).

The global species population is estimated at 15,337-30,519 pairs with a decreasing population trend (30% in the next 54 year i.e., three generations). Ten colonies in the Mediterranean Sea have disappeared during the last 60 years (Derhe 2012B, BirdLife International 2015, Birdlife International 2018B). The national population is estimated at 4,000-7,000 pairs (without the inclusion of the Gyaros colony which is estimated at 3,090-7,450 pairs), equivalent to 22% percent of the global population (more than 38% with the inclusion of the Gyaros population). The national population trend is estimated to be stable.

2.2.4.3 *European Storm-petrel (Hydrobates pelagicus)*



Figure 2-19. *European Storm-petrel (Hydrobates pelagicus)* (© Paul Hirst)

European Storm-petrel is the smallest seabird species in the Western Palaearctic. Its distribution is limited mainly to the Northeast Atlantic Ocean and the West Mediterranean Sea, while the Aegean Sea comprises the easternmost part of its range. The Mediterranean

subspecies *Hydrobates pelagicus melitensis* comprises less than 5% of the overall global population (i.e., 12,000-17,500 breeding pairs) with the main colonies located in Malta, Sicily and the Balearic Islands. The species occurs in all Greek seas mainly in spring and summer during the breeding period. Up to date only two colonies have been located, one in the Central Aegean Sea and another in the Cyclades. Storm-petrels, usually individual birds, or very small groups, are regularly observed in the Cyclades, Dodecanese, Central and southwest Aegean Sea and the Karpathanian Sea suggesting potential existence of other breeding colonies (Fric et al. 2012).

2.2.4.4 Mediterranean Shag (*Phalacrocorax aristotelis desmarestii*)



Figure 2-20. Mediterranean Shag (*Phalacrocorax aristotelis desmarestii*) (© Jens Overgaard Christensen)

Mediterranean Shag is a cormorant species, resident and widely spread in Greece which usually occurs in coastal waters. Shags breed colonially, forming small, loose (rarely dense) colonies, on cliff ledges or small caves or even under thick vegetation. Nesting sites are re-used in successive years by the same birds. They often roost in large groups (Fric et al. 2012). It is a good swimmer and a foot-propelled diver which feed on benthic and pelagic fish in waters with depths up to 80 m which are usually located in coastal zones within a 20 km radius around their colony or roosting sites (Wanless *et al.* 1991; Velando and Friere 1999).

The Greek national population size is 1,300 -1,450 pairs (Fric et al. 2012), equivalent to 2% of the species European population (BirdLife International 2015, BirdLife International 2018D). The population in Greece is considered to be stable (Fric et al. 2012). The island of Zakynthos hosts an important population of the Mediterranean Shag (i.e., 44-46 adult and juvenile individuals) (Portolou et al. 2009, Fric et al. 2012). The species breeds along the western coast of the Zakynthos and forages in coastal waters along the western and eastern coast, including the Bay of Laganas. Therefore, the major breeding and foraging areas of the Mediterranean Shag around Zakynthos Island are located outside the Project Area and within the Wider Project Area.

2.2.4.5 Yellow-legged Gull (*Larus michahellis*)




Figure 2-21. Yellow-legged Gull (*Larus michahellis*) (© Paul Hirst)

The Yellow-legged Gull is the most common gull species in Greece. It is widely distributed around the southern regions of the Palaearctic, from the western part of the Black Sea across to the Mediterranean, Iberian Peninsula, and reaching the Macaronesian region. Breeding grounds are centred mainly around the Mediterranean but reach also the Black Sea, Caspian Sea and eastern Atlantic. In Greece, the species is resident and widespread all along the coastline of mainland Greece and of the islands of the Aegean and Ionian Seas.

In Greece, the largest breeding colonies are located on uninhabited islets of the Evvoikos and Saronikos Gulfs that surround Attica, the most urbanised area in the country, although colonies occur on most Greek islets (Fric et al. 2012). Wintering grounds include the coast of southwest Asia, most of the European coast up to Denmark and the coast of Africa from Western Sahara through the eastern Mediterranean (del Hoyo et al. 1996).

3 Methodology

3.1 Pelagic surveys

Pelagic surveys for cetaceans, sea turtles and seabirds are carried out using i) a 15m sailing boat, ii) a 7,5m RIB boat and iii) a high wing, ultralight aircraft.

3.1.1 Boat surveys

Visual-based surveys

The method applied for visual surveying seabirds, cetaceans and sea turtles in the Pelagic surveys area is the **European Seabirds at Sea (ESAS)**, based on Tasker *et.al* 1984 and Champhuysen & Garthe 2004 and adopted to Greek/Mediterranean conditions through the LIFE-Nature project for the Identification of Marine Important Bird Areas (marine IBAs) in Greece, entitled “Concrete Conservation Actions for the Mediterranean Shag and Audouin’s Gull in Greece, including the Inventory of Relevant Marine IBAs”, LIFE07 NAT/GR/000285, (<http://www.ornithologiki.gr/en/seabirds>), as described in Fric & Gaganis 2009.

In summary, the method is aiming at systematically recording seabirds, cetaceans and sea turtles as well as human activities in the survey area, in transects by trained observers, from a boat which is moving at a constant low speed (<15 knots). Swimming seabirds, cetacean, fish and sea turtles are being recorded continuously in a 300m wide strip transect in **5-minute intervals**, while flying birds are recorded with **1-min snapshot**. Scanning angle is 180° (*i.e. in front of the survey vessel*). The perpendicular distance of swimming fauna is recorded relative to the transect line ahead of the ship: **A = 0-50m, B = 50-100m, C = 100-200m, D = 200-300m, E = >300m, W = within 300m, but no distance recorded**. For flying birds, coded with **F**, there is no distance indication. Boat position (**poskey**), namely geographical longitude and latitude, are recorded every 5 min. The marine species are spotted by a naked eye or binoculars and are identified by binoculars.

A method described by Heinemann (1981) is used to determine the distances at sea and more particularly the distance of 300m from the observing platform which determines the width of the line transect by using a calliper or a ruler. During ESAS surveys data is recorded regarding (A) boat route, (B) marine species and (C) human activities in the survey area, which may have an effect on the presence and behaviour of the marine species.

Survey boat data include: start and end location date, time and geographical location of each line transect, sea state, visibility and floating matter (including fishing vessels). Species data recorded include: species, number of individuals, age (if applicable), distance from the observation vessel, location within or outside 300m line transect, flight direction (for birds), behavior and association with human activities or other species. Datasheets for observation vessel data and species data are provided in Annex I.




Figure 3-1. ESAS field work

The **survey design for cetaceans** is similar to the established methodology designs for such surveys, used over the past 4 decades (Buckland et al. 2001, Buckland et al. 2004) and used a grid of parallel line transects, that provided comprehensive coverage of the study area.

The transect lines acted as the basis for the daily track line followed by the vessel providing a roughly uniform coverage of the study area. Attempts were made when selecting the orientation of the transect lines, to have them move across (at an angle to) the depth gradient in the area as opposed to moving along (parallel to) the depth gradient. This was done to allow for the coverage of different depth levels during navigation of each transect, in order to minimize detection bias on individual transect lines when mapping sighting data.

When a group of cetaceans is sighted (group defined 'dolphins observed in apparent association, moving in the same direction and often, but not always, engaged in the same activity' (Bearzi et al. 2005) by any of the on-effort observers, the systematic search effort is interrupted while the vessel diverted from the track line toward the sighted animals in order to achieve more accurate determinations of the species, the group size, group age class composition and group activity of the group sighted. In addition to basic environmental data (e.g., Beaufort sea state, visibility conditions etc.) collected at regular 1 hour intervals as well as at the start and at the end of each transect line, data collected for each sighting includes the time, GPS coordinates, initial bearing and radial distance to the cetacean group (used to calculate the perpendicular distance of the sighting to the track line), species identity, group size, group age class composition (3 age classes: Calf < 1/2 length of adult, Juvenile < 2/3 length of adult and adult) and the general activity in which the group is engaged in at the time of approach (e.g. foraging, travelling, milling). For the purpose of the correct identification of the species as well as the correct recording of group size and group age class composition

attempts are made to approach the animals to obtain photographs. Where possible the photographs taken are also used for the photo-identification of individuals. This is done to ensure the same group of animals was not counted twice during the same survey day.

Encounter Rates are calculated as the number of encounters / 100km of “on effort” navigation.

The navigation schedule coincided with the Visual boat-based surveys.

In case a group of cetaceans or seabirds was spotted, a drone was used in order to more accurately identify the species and assess the number of the individuals, record their behaviour and gather the relevant photographic evidence.

The numbers of individuals of each species recorded by ESAS surveys were transformed into species densities per km², taking into account the $2 \times 300m = 600m$ transect survey width and the distance travelled by the survey vessels per 5-minute time interval *distance travelled* = *boat speed* x 5 min. The locations of number of recorded individuals per species and the density of individuals per species were overlaid 4 geographical minutes (4'x4') reference grid in WGS84 projection coordinate system (Map 5).

Taking into account that more than one may have crossed each 4'x4' reference grid cell, for each cell the following variables were calculated:

- The **average** over all survey trips of the **total number of individuals per species** recorded in a 4'x4' grid cell per trip
- The **maximum** over all survey trips of the **total number of individuals per species** recorded in a 4'x4' grid cell per trip
- The **average** over all survey trips of the **average density of individuals per km² per species** in a 4'x4' grid cell per trip
- The **average** over all survey trips of the **maximum density of individuals per km² per species** in a 4'x4' grid cell per trip
- The **maximum** over all survey trips of the **average density of individuals per km² per species** in a 4'x4' grid cell per trip
- The **maximum** over all survey trips of the **maximum density of individuals per km² per species** in a 4'x4' grid cell per trip

It should be noted that individuals recorded outside transect are excluded from density calculation. The densities of the species per reference grid cell are representative of the **habitat suitability**. The variable “**average** over all survey trips of the **average density of individuals per km² per species** in a 4'x4' grid cell per trip” was used as a measure of habitat suitability for each species. This variable was classified into 4 classes:

- **Most suitable habitats** – top 5% of positive (i.e., non-zero) densities in grid cells
- **More suitable habitats** – 25-5% top values of positive densities in grid cells

- **Suitable habitats** – 75-25% top values of positive densities in grid cells, and
- **Presence** – remaining grid cells with species presence (bottom 25% values).

To further analyse the **patterns of seabird movements** in the area for each grid cell the **prevailing flight directions** were calculated. Additionally, **locations of interactions of seabirds with fisheries** were identified in association with their abundance in absolute numbers.

Finally, for each grid cell the **number of species of interest recorded** in the grid cell was calculated to identify those areas where the **species richness** is the greatest.




Figure 3-2. Four decimal minute (4'x4') reference grid in WGS84 coordinate system.

Acoustic surveys

The acoustics detection team works in cooperation with the visual observers, detecting cetacean vocalizations by using a hydrophone array towed behind a sailing boat. The hydrophone array system consisting of High Frequency Magrec HP03 hydrophone elements, comprising a HP03 preamp (Low cut filter set at 2kHz) with a nominal sensitivity of 1.5kHz – 150kHz along with a topside Magrec HP/27ST Amplifier along with a Lenovo Thinkpad Laptop using the PAMGUARD acoustic analysis software specifically developed for cetacean monitoring, covering the range of possible vocalizations for species likely to be encountered during our surveys. The towed hydrophone system is submerged and active, and a PAM operator is active on the equipment during all “On Effort” times during the survey. The hydrophone system consists of 2 hydrophones which record in 2 different channels. The visual observers and PAM operator rotate every 1.5 hour to minimize fatigue.

The PAM operator immediately informs the visual observer team of any acoustic detection.

The hydrophone recordings are analysed by PAMGUARD software using “*whistle and moan detector*” module.



Figure 3-3. Part of the hydrophone array towed behind a sailing boat

3.1.2 Aerial surveys

For the aerial surveys, a high wing, light aircraft Cessna C172R Skyhawk was used, powered with a Lycoming IO-360-L2A, 160 Hp. This four-seater aircraft offers an excellent view from its cockpit and is considered suitable, reliable and cost-effective for such a mission. Messolonghi Airport (ICAO designator GR-0008) was used as a base for the aerial expeditions to the Northern Ionian Project Area. The flight was performed along the Project Area at an altitude of 1000 ft MSL and an average Speed Over Ground of 85 knots. The flights were performed under ideal weather conditions (wind speed less than 10 knots, clear sky and

visibility more than 8 km). In every case where an “object/s of interest” was spotted, the airplane left its track and performed one or more circles over the object/s in order to visually identify it. Furthermore, the object was photographed so that a proper record of its observation and identification is kept. The photographic operation was performed using a full frame DSLR (Nikon D750) with a 70-200mm F/2.8 Tamron SP lens. All photographs were georeferenced since the camera was equipped with a GPS Unit (Nikon GP-1A). The flights were monitored and recorded with the use of two specialized applications namely the GARMIN-Pilot and the Fore-Flight.



Figure 3-4. The aircraft used for the aerial surveys



Figure 3-5. View from the aircraft's cockpit

In the following example, the staged photographic identification process of an initially “object of interest” located on the shore is clearly shown.



Figure 3-6. A: Recording an “object of interest”, B: Approaching, C: Identifying

3.2 Coastal surveys

3.2.1 Coastal surveys for the Scopoli's Shearwater

At species' breeding colony in Strofadia islands, an adaptation of the existing raft counting method has been developed, with the RIB boat following the birds gathering in front of the colony before sunset, to create the raft. When a raft is spotted, the number of birds is counted using binoculars and ZOOM cameras. The DJI Mini 2 drone is then deployed flying at 30m above sea level to take photos and 4k video of the raft, in order to provide more accurate estimations.

At a second stage, after sunset, the raft is further monitored using a 640x480 thermal camera, to assess the movements of birds from the rafts to the colonies, as well as the timeline of the birds entrance and flights to the colony sites. In this respect, breeding birds are distinguished from prospectors to provide more precise estimates of the colony size.

3.2.2 Coastal surveys for the Mediterranean Shag

Coastal surveys for the Mediterranean Shag involve the recording of the species individuals, age and activity while the survey vessel travels at a low speed along the survey coastline at a distance of 50-100m from the shore. The species are identified by binoculars, data is recorded on field maps and their locations are recorded by a portable GPS unit. Simultaneously, apparently active or suspected nesting sites are recorded.

The data recorded during field surveys included:

- Date / time of the observation
- Location of the observation (GPS waypoint name, latitude, longitude)
- Seabird species
- Number of individuals
- Number of adult and juvenile individuals (for the Mediterranean Shag)
- Identification of colony/nest sites, number of nests, suitable nesting habitat, roosting sites
- Potential localised threats
- Comments

3.2.3 Coastal surveys for the Mediterranean Monk Seal

Cave monitoring with the use of Automated Infrared Cameras

During the previous phase of the project, 4 Infrared Camera Traps were placed in 2 caves in the wider project area: ZAK6 and ZAK7. Three camera systems were installed in cave ZAK6 and 1 was installed in cave ZAK7. These were placed on rocky walls of the caves and were positioned to maximize the coverage of the terrestrial component of the cave (internal beach). Three of the camera systems were set to take time lapse photos (every 1 and 2.5 hours) of the cave beach to record the presence or absence of animals while one camera system set on motion detection operation.



Figure 3-7. Installation of infrared camera systems in a monk seal shelter in the project area

3.3 Sea turtle telemetry

The fieldwork therefore had two main goals:

- 1) Retrieve transmitters from male turtles tracked from 2023 to access stored data.
- 2) Deploy transmitters on a further two male turtles

To identify if the turtles from 2023 had returned to Kyparissia in 2024 we deployed a drone (DJI Phantom 3) to systematically search the field site for evidence of turtles with transmitters. This was undertaken throughout the field period, whilst the other work was being carried out.

We acquired the two male turtles to receive transmitters using turtle rodeo technique as used in the previous fieldwork, namely a small RIB was used to approach the turtles and then they were caught by hand, secured in a net and taken ashore to receive the transmitters. Should we have seen a male turtle with a transmitter in place, we would have used the same rodeo technique to catch it to retrieve the transmitter.



Figure 3-8. Wildlife Computers SPLASH10-385 satellite transmitter used in the project.




Figure 3-9. Left to right: The RIB and drone used to spot and catch the turtles. Capturing the turtle ready to bring ashore. Tagging and other field equipment used to record details of the captured turtle.

3.4 Telemetry of Scopoli's Shearwater

The bird flights at the species breeding colony at Strofades islands will be recorded by a drone and stationary thermal cameras and will be stored to a geodatabase.

The birds movements will be monitored in the vicinity of island colonies and foraging aggregations of seabirds and marine mammals will be spotted and mapped within the project area. An array of trail cameras and bio-acoustic microphones will be deployed at the Strofades colony to record the colony activities, threats and social interactions among breeding and non-breeding individuals.

10 GPS/GSM tags will be deployed to Scopoli's shearwater fledglings on September-early October 2024, to record the maiden journeys of the fledged birds and identify possible threats and mortality factors during this high-risk period.

By processing the data through Artificial Intelligence and machine learning software, the bird populations, movements, patterns of space use by the species for foraging in the "Kyparissiakos block" lease area will be further explored.

4 Results

4.1 Pelagic surveys

4.1.1 Boat surveys

A total of **964 nautical miles** of boat-based visual surveys were carried out on 4-7/7/2024, 13-19/7/2024 and 7-11/10/2024 in the Pelagic Survey Area, as well as in the surrounding areas in the Wider Project Area, to assess the presence, abundance and distribution of the cetacean, sea turtle and seabird species of interest.




Figure 4-1. Visual boat-based survey tracks

During the visual surveys the following species were recorded:

- 3 cetacean species: **Sperm whale** (*Physeter macrocephalus*), **Cuvier's beaked whale** (*Ziphius cavirostris*), and **stripped dolphin** (*Stenella coeruleoalba*),
- 6 seabird species: **Scopoli's Shearwater** (*Calonectris diomedea*), **Yelkouan Shearwater** (*Puffinus yelkouan*), **Yellow-legged Gull** (*Larus michahellis*), **Mediterranean Shag** (*Phalacrocorax aristotelis*), **European Storm Petrel** (*Hydrobates pelagicus*) and **Common Tern** (*Sterna hirundo*),
- 1 wetland bird species: **Shelduck** (*Tadorna tadorna*),
- 1 raptor species: **Marsh Harrier** (*Circus aeruginosus*),
- 1 sea turtle species: **Loggerhead turtle** (*Caretta caretta*).

Table 4-1. Species recorded in the Project Area

Species	Common name	Number of individuals
<i>Physeter macrocephalus</i>	Sperm whale	3
<i>Ziphius cavirostris</i>	Cuvier's beaked whale	9
<i>Stenella coeruleoalba</i>	Bottlenose dolphin	2
Dolphin (unidentified species)		3
<i>Calonectris diomedea</i>	Scopoli's Shearwater	335
<i>Puffinus yelkouan</i>	Yelkouan Shearwater	100
<i>Hydrobates pelagicus</i>	European Storm Petrel	1
<i>Larus michahellis</i>	Yellow-legged Gull	35
<i>Sterna hirundo</i>	Common Tern	1
<i>Phalacrocorax aristotelis</i>	Mediterranean Shag	3
<i>Tadorna tadorna</i>	Shelduck	12
<i>Circus aeruginosus</i>	Marsh Harrier	1
<i>Caretta caretta</i>	Loggerhead turtle	1

Moreover, during the visual surveys, the following cetacean species were observed outside the Wider Project Area:

- **3 sperm whales** NW of Zakynthos Island,
- **1 sperm whale** in the deep waters of South Messiniakos Gulf,
- **4 Cuvier's beaked whales**, W of Zakynthos Island,
- **28 striped dolphins**, W of Kefalonia Island.

During the **acoustic surveys with the towed hydrophone**, covering 827 nautical miles, **36 detections of sperm whales** were recorded, as well as **16 detections of dolphins** (species identification is not possible with the towed hydrophone survey) in the Wider Project Area.

Additionally, another **21 acoustic detections of sperm whales** were recorded outside the Wider Project Area (NW of Zakynthos and W of Kefalonia Islands).

4.1.2 Aerial surveys

The aerial survey was conducted on the 7th of April 2024. A total of **644 km** of coastline were inspected, covering the project area, as well as the western coast of Zakynthos Island and the northwestern coast of Peloponnese.




Figure 4-2. Aerial surveys track.

Cetaceans




Figure 4-3. Locations of sperm whale visual records.




Figure 4-4. Locations of Cuvier's beaked whale visual records.




Figure 4-5. Locations of striped dolphin and unidentified dolphins visual records.

Seabirds




Figure 4-6. Locations of Scopoli's Shearwater visual records.




Figure 4-7. Locations of Yelkouan Shearwater visual records.




Figure 4-8. Locations of Storm Petrel visual records.







Figure 4-9. Sperm whales during the boat surveys.



Figure 4-10. Cuvier's Beaked whale (above) and striped dolphins (below) during the boat surveys.



Figure 4-11. Sperm whale with ferry and sailing boats(above) and Scopoli's Shearwater (below).



Figure 4-12. A group of 4 grey herons (Ardea Cinerea) migrating in the project area during the aerial surveys.



Figure 4-13. A trawler operating in the project area during the aerial surveys.

4.2 Coastal surveys

4.2.1 Coastal surveys for the Scopoli's shearwater

Coastal surveys for Scopoli's shearwater were conducted in July and September 2024 using a RIB boat. The coastal surveys focused on Strofades islands, where the largest breeding colony of Scopoli's Shearwater in Greece is located. Both drone surveys and thermal camera monitoring took place, as well as a visit to the colony on the islet of Stamfani to locate accessible nests for tagging birds.

4.2.2 Coastal surveys for the Mediterranean Shag

Coastal surveys for the Mediterranean shag at the Kyparissia coastline and the Strofades islets revealed no birds. This is attributed to the very low densities of the species in the project area, due to the unsuitable habitat (Kyparissia) or the offshore character of the project sites.

4.2.3 Coastal surveys for the Mediterranean Monk Seal

During the previous stages of the project, the coastline within the Wider Project Area was surveyed for the presence of suitable monk seal pupping habitat (marine caves). Monitoring efforts of the monk seal population during the previous reporting period focused on the island of Zakynthos. Three automatic infrared camera systems were placed in 2 pupping caves selected in the island of Zakynthos: ZAK6 and ZAK7. As Zakynthos island's caves and the monk seal population using them are the most vulnerable in relation to any future drilling activities and potential spills in the Kyparisiakos lease area, monitoring of these caves was considered essential.

The results of the infrared cameras in combination with data collected through the operation of the Rescue and Information Network (RINT) of MOM in the area are summarized up as follows:

- 21 monk seal observations collected through RINT
- 261 days of cave camera monitoring conducted
- 13329 photographs collected
- 17 different individuals identified through the analysis of the photos captured:
 - 7 Adult females
 - 2 Adult males
 - 2 Sub-adults
 - 1 Female pup
 - 3 Male pups
 - 2 pups of unknown sex

Based on the above-mentioned information, the number of monk seals inhabiting the area of Zakynthos Island is estimated to be at least 20 individuals

These results documented for the first time the importance of Zakynthos Island as a monk seal pupping site in the Ionian region.



Figure 4-14. The earliest-born pup in cave ZAK6 suckling from its mother along two other adult females and a sub-adult.



Figure 4-15. Animals of all age classes photographed in ZAK6.

Please refer to Annex I for the full report on cave monitoring and demographic parameters of the Mediterranean Monk Seal.

4.3 Sea turtle telemetry

Despite multiple hours of searching with the drone, with many turtles observed, none was recognised as having a transmitter in place. We therefore concluded that the two previously tracked turtles had not returned to Kyparissia in 2024. Our survey methods were validated as the drone observed one of the 2024 turtles in the field area after it had received its transmitter.



Figure 4-16. Left: Three turtles observed in close proximity to each other during drone surveys. None have a transmitter in place. Right: The first turtle to receive a transmitter in 2024 observed during the drone surveys after his release.

On the deployment side, we were able to successfully catch the planned two adult male turtles and equip them with transmitters. Both turtles' transmitters continue to transmit locations and data and they were recorded making their post-breeding migrations to widely disparate foraging locations.






Figure 4-17. The two adult males that were successfully equipped with satellite transmitters during the April 2024 field mission.





Please refer to Annex II for the full report on Loggerhead turtles tracking in Kyparissia.

4.4 Telemetry of Scopoli's Shearwater

The movements and flight patterns of Scopoli's Shearwater were monitored by telemetry. The birds were tagged at the breeding colony of the species on Strofades Islands. It is worth mentioning that for the first time in Greece, both adults and young birds were tagged, giving us the opportunity to study not only the movements of the parents during the breeding season, but also the dispersal patterns of the young birds after leaving their nests.

The results revealed that the tagged adult birds used the south and central Ionian Sea and Adriatic Sea, as well as the Gulf of Taranto and the eastern coast of Sicily, for foraging during their breeding season. Additionally, an individual regularly visited Gulf of Corinth for foraging. After the end of the breeding season, both adults and young birds left the colony on Strofades and headed towards the Western Mediterranean to reach their wintering grounds in the Atlantic.



Figure 4-19. Flight movements of the tagged Scopoli's Shearwaters.

5 Conclusions

During 2024, all the scheduled tasks have been performed successfully.

Both boat surveys (visual-based and acoustic) and aerial surveys were carried out in the project area in spring, summer and autumn 2024. In total 3 species of cetaceans (sperm whale, Cuvier's beaked whale and striped dolphin) were spotted, all typical cetacean species of the deep waters of the Project Area. The record of 9 Cuvier's beaked whales was significant, but the most important observation was the groups of sperm whales, both within and outside the Project Area. It is worth mentioning that this year a dedicated effort to locate sperm whales was carried out under the guidance of Dr. Jonathan Gordon, an international expert of the species, who supported the overall fieldwork project planning this year and participated in the surveys in July. These observations confirm that the Hellenic Trench in Ionian Sea, at depths >1000m, is the most suitable marine habitat for the large cetaceans of Greece.

Additionally, 6 species of seabirds and one species of sea turtle were recorded during the surveys. Scopoli's Shearwater was the most abundant seabird species, as it holds the largest breeding colony of Greece within the Project Area, on Strofades islands, and regularly uses the surrounding marine area as a foraging ground.

Telemetry of Scopoli's Shearwater revealed that the tagged birds used the wider area of North Ionian – South Adriatic Sea as foraging grounds during their breeding period. During the first weeks of October, both adults and young birds left the colony and started moving towards the Western Mediterranean in order to arrive at their wintering grounds in Atlantic.

Regarding sea turtle telemetry, transmitters were deployed on two adult male Loggerhead turtles in Kyparissia. Telemetry of adult males is providing novel data on their behaviour and movements, helping fill a gap in the understanding of sea turtle biology, not only in Greece but globally. By the end of October 2024, one male has been moved to the eastern part of Gulf of Corinth, while the other headed southwest towards the Gulf of Gabes.

Regarding the Mediterranean Monk Seal, the results of the current Monitoring Survey for the species population of Zakynthos Island provide an updated (and for the first time well documented) assessment of the status of this rare species in the area.

The evidence collected during the project verify Zakynthos Island as indeed being a highly important hub for the Mediterranean monk, not only in the Kyparissiakos Gulf wider area but in the Ionian Sea in general. Specifically cave ZAK6 shows an intense use by monk seals to give birth and raise their pups.

The up-to-date findings confirm the necessity to establish prevention and response measures and contingency plans for the protection of the pupping habitat of the monk seals on the island in the event of an oil spill accident during drilling operations in the region.

Bibliography

- Adamantopoulou, S., Androukaki, E. & Kotomatas, S. (1999). The distribution of the Mediterranean monk seal in Greece based on an information network. *Contributions to the Zoogeography and Ecology of the Eastern Mediterranean Region*, 1, 399-404
- Aguilar, A. & Gaspari, S. 2012. *Stenella coeruleoalba*. The IUCN Red List of Threatened Species 2012: e.T20731A2773889.
- Aguilar, A. & Lowry, L. (IUCN SSC Pinniped Specialist Group). 2010. *Monachus monachus*. The IUCN Red List of Threatened Species 2010: e.T13653A4305567.
- Anonymous (2007). Status of the population of the Mediterranean monk seal (*Monachus monachus*) in Greece. pp. 1-42. MOM/Hellenic Society for the Study and Protection of the Monk seal, Athens, Greece.
- Avens, L. and M.L. Snover (2013) Age and age estimation in sea turtles. In: J. Wyneken, K.J. Lohmann & J.A. Musick (ed.), *The biology of sea turtles. Volume III.*, pp. 97-133. CRC Press, Boca Raton, FL, USA.
- Bearzi, G. 2003. *Delphinus delphis* (Mediterranean subpopulation). The IUCN Red List of Threatened Species 2003: e.T41762A10557372.
- Bearzi, G., Reeves, R.R., Notarbartolo di Sciara, G., Politi, E., Cañadas, A., Frantzis, A. and Mussi, B. 2003. Ecology, status and conservation of Short-beaked Common Dolphins (*Delphinus delphis*) in the Mediterranean Sea. *Mammal Review*. 33(34): 224-252.
- Bearzi, G., Politi, E., Agazzi, S., Bruno, S., Costa, M. and S. Bonizzoni (2005) Occurrence and present status of coastal dolphins (*Delphinus delphis* and *Tursiops truncatus*) in the eastern Ionian Sea. *Aquatic Conserv: Mar. Freshw. Ecosyst*. 15: 243–257
- Bearzi, G., Fortuna, C. M. and R. R. Reeves (2008A) Ecology and conservation of common bottlenose dolphins *Tursiops truncatus* in the Mediterranean Sea. *Mammal Rev*. 39(2): 92-123.
- Bearzi, G., Agazzi, S., Gonzalvo, J., Costa, M., Bonizzoni, S., Politi, E., Pirroddi, C. and R.R. Reeves (2008B) Overfishing and the disappearance of short-beaked common dolphins from western Greece. *Endang. Species. Res*. 5: 1-12
- Bearzi, G., Fortuna, C. & Reeves, R. 2012. *Tursiops truncatus*. The IUCN Red List of Threatened Species 2012: e.T22563A2782611.
- BirdLife International (2015) European Red List of Birds. Luxembourg: Office for Official Publications of the European Communities
- BirdLife International (2018A) *Calonectris diomedea*. The IUCN Red List of Threatened Species 2018: e.T45061132A132667885. Downloaded on 15 November 2018.
- BirdLife International (2018B) *Puffinus yelkouan*. The IUCN Red List of Threatened Species 2018: e.T22698230A132637221. Downloaded on 15 November 2018.
- BirdLife International (2018C) *Hydrobates pelagicus*. The IUCN Red List of Threatened Species 2018: e.T22698477A132650209. Downloaded on 15 November 2018.

- BirdLife International (2018D) *Gulosus aristotelis*. *The IUCN Red List of Threatened Species* 2018: e.T22696894A133538524. Downloaded on 15 November 2018.
- BirdLife International (2018E) *Larus audouinii*. *The IUCN Red List of Threatened Species* 2018: e.T22694313A132541241. Downloaded on 15 November 2018.
- BirdLife International (2018F) *Larus michahellis*. *The IUCN Red List of Threatened Species* 2018: e.T62030970A132234755. Downloaded on 15 November 2018.
- Bolten, A.B. and B.E. Witherington (2003) *Loggerhead Sea Turtles*. Smithsonian Books, Washington, D.C., USA.
- Bourgeois, K. & E. Vidal (2008) The endemic Mediterranean yelkouan shearwater *Puffinus yelkouan*: distribution, threats and a plea for more data. *Oryx* 42(2): 187-194.
- Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L. and L. Thomas (2001) *Introduction to Distance Sampling: Estimating Abundance of Biological Populations*. Oxford University Press, Oxford, UK.
- Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L. and L. Thomas (2004) *Advanced Distance Sampling: Estimating Abundance of Biological Populations*. Oxford University Press, Oxford, UK.
- Cañadas, A. & Notarbartolo di Sciara, G. 2018. *Ziphius cavirostris* (Mediterranean subpopulation). *The IUCN Red List of Threatened Species* 2018: e.T16381144A50286386.
- Carboneras, C., Derhé, M. & I. Ramirez (2013) Update on the population status and distribution of Mediterranean shearwaters. Report to Seventh Meeting of the ACAP Advisory Committee, La Rochelle, France, 6-10 May 2013.
- Casale, P. & D. Margaritoulis (Eds.). (2010). *Sea turtles in the Mediterranean: distribution, threats and conservation priorities*. IUCN.
- Casale, P. & P. Mariani (2014) The first 'lost year' of Mediterranean sea turtles: dispersal patterns indicate subregional management units for conservation. *Mar Ecol Prog Ser* 498: 263–274
- Casale, P. (2015) *Caretta caretta Mediterranean subpopulation*. *The IUCN Red List of Threatened Species* 2015: e.T83644804A83646294.
- Casale, P. & A.D. Tucker (2017) *Caretta caretta* (amended version of 2015 assessment). *The IUCN Red List of Threatened Species* 2017: e.T3897A119333622.
- Casale, P., Broderick, A. C., Camiñas, J. A., Cardona, L., Carreras, C., Demetropoulos, A., Fuller, W. J., Brendan, J. G., Hochscheid, S., Kaska, Y., Lazar, B., Margaritoulis, D., Panagopoulou, A., Rees, A.F., Tomás, J. & Türkozan, O. (2018). Mediterranean sea turtles: current knowledge and priorities for conservation and research. *Endangered species research*, 36, 229-267.
- Champhuysen, C. J. & S. Garthe (2004) Recording foraging seabirds at sea: Standardized recording and coding of foraging behaviour. *Atlantic Seabirds* 5(si)

- Dendrinou, P., Karamanlidis, A.A., Kotomatas, S., Legakis, A., Tounta, E. & Matthiopoulos, J. (2007). Pupping habitat use in the Mediterranean monk seal: a long-term study. *Marine Mammal Science*, 23, 615-628.
- Dendrinou, P., Kotomatas, S. & Tounta, E. (1999). Monk seal pup production in the National Marine Park of Alonissos-N.Sporades. *Contributions to the Zoogeography and Ecology of the Eastern Mediterranean Region*, 1, 413-419.
- Derhé, M. A. (2012A) Developing a Population Assessment for Scopoli's and Cory's Shearwaters *Calonectris diomedea/Calonectris borealis*. In: Yésou, P.; Baccetti, N.; Sultana, J. (ed.), *Ecology and Conservation of Mediterranean Seabirds and other bird species under the Barcelona Convention – Proceedings of the 13th Medmaravis Pan- Mediterranean Symposium. Alghero (Sardinia) 14-17 Oct. 2011*, pp. 29–38. Medmaravis, Alghero
- Derhé, M. A. (2012B) Population assessment for the Yelkouan Shearwater *Puffinus yelkouan*. In: BirdLife International (ed.), *Methodology for Bird Species Recovery Planning in the European Union. Final Report to the European Commission*. BirdLife International for the European Commission, Cambridge, UK.
- Dimalaxis, T., Kastiris, Th., Grivas, K., Manolopoulos, A., Kardakari, N., Kakalis, L., Xirouchakis, S., Tsaitouridis, S., Papazoglou, C. and B. Barov (2009) Definition of compatible activities with respect to the species for the designation of the Special Protection Areas for birds. [In Greek]
- del Hoyo, J., Elliott, A. and J. Sargatal (1996) *Handbook of the Birds of the World*, vol. 3: Hoatzin to Auks. Lynx Edicions, Barcelona, Spain.
- Diogou, N., Klinck, H., Frantzis, A., Nystuen, J. A., Papathanassiou, E., & Katsanevakis, S. (2019). Year-round acoustic presence of sperm whales (*Physeter macrocephalus*) and baseline ambient ocean sound levels in the Greek Seas. *Mediterranean Marine Science*, 20(1), 208-221.
- Frantzis A., (1998) Does acoustic testing strand whales? *Nature* 392:29
- Frantzis A., Alexiadou P., Paximadis G., Politi E., Gannier A., Corsini-Foka M., 2003. Current knowledge of the cetacean fauna of the Greek Seas. *The Journal of Cetacean Research Management* 5(3): 219-232.
- Frantzis A. (2009) *Cetaceans in Greece: Present status of knowledge*. Initiative for the Conservation of Cetaceans in Greece, Athens, Greece, 94 pp.
- Frantzis, A., Alexiadou, P., & Gkikopoulou, K. C. (2014). Sperm whale occurrence, site fidelity and population structure along the Hellenic Trench (Greece, Mediterranean Sea). *Aquatic Conservation: Marine and Freshwater Ecosystems*, 24(S1), 83-102.
- Frantzis, A., Leaper, R., Alexiadou, P., Prospathopoulos, A., & Lekkas, D. (2019). Shipping routes through core habitat of endangered sperm whales along the Hellenic Trench, Greece: Can we reduce collision risks?. *PloS one*, 14(2), e0212016.
- Fric J. & K. Gaganis 2009. European Seabirds at Sea Methodology. LIFE-Natura project LIFE07 NAT/GR/000285, Hellenic Ornithological Society. [In Greek]

- Fric, J., Portolou, D., Manolopoulos, A. and T. Kastritis (2012). *Important Areas for Seabirds in Greece*. LIFE07 NAT/GR/000285 - Hellenic Ornithological Society (HOS / BirdLife Greece), Athens.
- Fric, J. & D. Portolou (2016). Population study of the Yelkouan Shearwater (*Puffinus yelkouan*) in the area of Gyaros. 2015 annual report. Nature Conservation Consultants (NCC) Ε.Π.Ε. & Hellenic Ornithological Society [In Greek].
- García-Barcelona, S., Ortiz de Urbina, J. M., de la Serna, J. M., Alot, E. & D. Macias (2010) Seabird bycatch in Spanish Mediterranean large pelagic longline fisheries, 2000-2008. *Aquatic Living Resources* 23: 363–371.
- Gaspari, S. & Natoli, A. 2012. *Grampus griseus* (Mediterranean subpopulation). The IUCN Red List of Threatened Species 2012: e.T16378423A16378453.
- González, L.M. & Fernandez de Larrinoa, P. (2012). Mediterranean monk seal *Monachus monachus* distribution and fisheries interactions in the Atlantic Sahara during the second half of the 20th century. *Mammalia*, 77, 41-49.
- Granadeiro, J. P., Dias, M. P., Rebelo, R., Santos, C.D. & P. Catry (2006) Numbers and population trends of Cory's Shearwater *Calonectris diomedea* at Selvagem Grande, Northeast Atlantic. *Waterbirds* 29: 56-60.
- Güçlüsoy, H., Kiraç, C.O., Veryeri, N.O. & Savaş, Y. (2004). Status of the Mediterranean monk seal, *Monachus monachus* (Hermann, 1779) in the coastal waters of Turkey. *E.U. Journal of Fisheries & Aquatic Sciences*, 21, 201-210.
- Gücü, A.C., Gücü, G. & Orek, H. (2004). Habitat use and preliminary demographic evaluation of the critically endangered Mediterranean monk seal (*Monachus monachus*) in the Cilician Basin (Eastern Mediterranean). *Biological Conservation*, 116, 417-431.
- Heinemann D. 1981. A range finder for pelagic bird censusing. *J. Wildl. Manage.* 45(2), pp. 489-493.
- Haywood, JC, Fuller WJ, Godley BJ, Magaritoulis D, Shutler JD, Snape RTE, Widdicombe S, Zbinden JA, Broderick AC (2020) Spatial ecology of loggerhead turtles: Insights from stable isotope markers and satellite telemetry. *Diversity and Distributions* 26: 368-381.
- IUCN (2012). *Marine Mammals and Sea Turtles of the Mediterranean and Black Seas*. Gland, Switzerland and Malaga, Spain: IUCN. 32 pages.
- Karamanlidis, A.A., Dendrinos, P., Fernández de Larrinoa, P., Gücü, A.C., Johnson, W.M., Kiraç, C.O. & Pires, R. (2016). The Mediterranean monk seal *Monachus monachus*: status, biology, threats, and conservation priorities. *Mammal Review*, 46, 92-105.
- Karamanlidis, A. & Dendrinos, P. 2015. *Monachus monachus* (errata version published in 2017). The IUCN Red List of Threatened Species 2015: e.T13653A117647375
- Karamanlidis, A.A., Pires, R., Silva, N.C. & Neves, H.C. (2004). The availability of resting and pupping habitat for the critically endangered Mediterranean monk seal *Monachus monachus* in the archipelago of Madeira. *Oryx*, 38, 180-185.
- Karris, G., Xirouchakis, S., Grivas, K., Fric, J., Dimalexis, T. and S. Sfenthourakis (2011) Migratory behaviour of Cory's Shearwaters, *Calonectris diomedea*, from an Ionian Sea colony: An

- application of miniature geolocation technology. In: Fusani L., Coppack T. and M. Stradzs (eds.) *Programme and Abstracts of the 8th Conference of the European Ornithologists' Union*, 27-30 August 2011, Riga, Latvia. Latvian Ornithological Society, 432 pp.
- Karris, G., Xirouchakis, S., Grivas, C., Voulgaris, M. D., Sfenthourakis, S., & S. Giokas (2017). Estimating the population size of Scopoli's Shearwaters (*Calonectris diomedea*) frequenting the Strofades islands (Ionian Sea, western Greece) by raft counts and surveys of breeding pairs. *North-Western Journal of Zoology*, 13(1).
- Karris, G., Xirouchakis, S., Maina, I., Grivas, K., & Kavadas, S. (2018). Home range and foraging habitat preference of Scopoli's shearwater *Calonectris diomedea* during the early chick-rearing phase in the eastern Mediterranean. *Wildlife Biology*, 2018(1).
- Laneri, K., Louzao, M., Martínez-Abraín, A., Arcos, J.M., Belda, E., Guallart, J., Sánchez, A., Giménez, M., Maestre, R. & D. Oro (2010). Trawling moratoria influences longline seabird bycatch in the Mediterranean: new insights from a small scale fishery. *Marine Ecology Progress Series* 420: 241–252.
- Legakis, A. and P. Maragou (2009) The Red Book of endangered animals of Greece. Athens, GR : Hellenic Zoological Society. 525pp
- Lewis, T., Gillespie, D., Lacey, C., Matthews, J., Danbolt, M., Leaper, R., McLanaghan, R. & Moscrop, A. (2007). Sperm whale abundance estimates from acoustic surveys of the Ionian Sea and Straits of Sicily in 2003. *Journal of the Marine Biological Association of the United Kingdom*, 87(1), 353-357.
- Luschi, P., & Casale, P. (2014). Movement patterns of marine turtles in the Mediterranean Sea: a review. *Italian Journal of Zoology*, 81(4), 478-495.
- Mannocci, L., Roberts, J. J., Halpin, P. N., Authier, M., Boisseau, O., Bradai, M. N., ... & Fortuna, C. M. (2018). Assessing cetacean surveys throughout the Mediterranean Sea: a gap analysis in environmental space. *Scientific Reports*, 8(1), 1-14.
- Margaritoulis, D. and A. Rees (2001) The Loggerhead Turtle, *Caretta caretta*, population nesting in Kyparissia Bay, Peloponnesus, Greece: results of beach surveys over seventeen seasons and determination of the core nesting habitat. *Zoology in the Middle East* 24: 75-90.
- Margaritoulis, D. (2005) Nesting activity and reproductive output of loggerhead sea turtles, *Caretta caretta*, over 19 seasons (1984-2002) at Laganas Bay, Zakynthos, Greece: the largest rookery in the Mediterranean. *Chelonian Conservation and Biology* 4: 916-929.
- Martínez-Jauregui, M., Tavecchia, G., Cedenilla, M.A., Coulson, T., Fernández de Larrinoa, P., Muñoz, M. & González, L.M. (2012). Population resilience of the Mediterranean monk seal *Monachus monachus* at Cabo Blanco peninsula. *Marine Ecology Progress Series*, 461, 273-281.
- Mo, G., Bazairi, H., Bayed, A. & Agnesi, S. (2011). Survey on Mediterranean monk seal (*Monachus monachus*) sightings in Mediterranean Morocco. *Aquatic Mammals*, 37, 248-255.
- Notarbartolo di Sciara, G. and J. Gordon (1997) Bioacoustics: A tool for the conservation of cetaceans in the Mediterranean Sea. *Marine & Freshwater Behaviour & Phy.* 30: 125-146

Notarbartolo di Sciara, G., Frantzis, A., Bearzi, G. & Reeves, R. 2012. *Physeter macrocephalus* (Mediterranean subpopulation). The IUCN Red List of Threatened Species 2012: e.T16370739A16370477.

Oppel S., Raine A.F., Borg J.J., Raine H., Bonnaud E., Bourgeois K. & A.R. Breton (2011) Is the Yelkouan shearwater *Puffinus yelkouan* threatened by low adult survival probabilities? *Biological Conservation* 144(9): 2255-2263.

Pires, R., Neves, H.C. & Karamanlidis, A.A. (2008). The Critically Endangered Mediterranean monk seal *Monachus monachus* in the archipelago of Madeira: priorities for conservation. *Oryx*, 42, 278-285.

Podestà, M., Azzellino, A., Cañadas, A., Frantzis, A., Moulins, A., Rosso, M., Tepsich, P. & Lanfredi, C. (2016). Cuvier's Beaked Whale, *Ziphius cavirostris*, distribution and occurrence in the Mediterranean Sea: high-use areas and conservation threats. In *Advances in Marine Biology* (Vol. 75, pp. 103-140). Academic Press.

Portolou, D., Bourdakis, S., Vlachos, C., Kastritis, T. and T. Dimalexis (eds.) (2009). *Important Bird Areas of Greece: Priority sites for conservation*. Hellenic Ornithological Society, Athens. [In Greek].

Rees, A. F., Theodorou, P., & Margaritoulis, D. (2020). Clutch frequency for Loggerhead Turtles (*Caretta caretta*) nesting in Kyparissia Bay, Greece. *Herpetological Conservation and Biology*, 15(1), 131-138.

Reeves, R. R., & Notarbartolo di Sciara, G. (2006). The status and distribution of cetaceans in the Black Sea and Mediterranean Sea.

Ristow, D., Berthold, P., Hashmi, D. and U. Querner (2000) Satellite tracking of Cory's shearwater migration. *Condor* 102: 696-699.

Santostasi, N. L., Bonizzoni, S., Gimenez, O., Eddy, L., & Bearzi, G. (2018). Common dolphins in the Gulf of Corinth are Critically Endangered. *Aquatic Conservation: Marine and Freshwater Ecosystems*.

Schofield, G., Hobson, V.J., Fossette, S., Lilley, M.K.S., Katselidis, K.A. & Hays, G.C. (2010a) Fidelity to foraging sites, consistency of migration routes and habitat modulation of home range by sea turtles. *Diversity & Distributions*, 16, 840–853.

Schofield, G., Hobson, V.J., Lilley, M.K.S., Katselidis, K.A., Bishop, C.M., Brown, P. & Hays, G.C. (2010b) Interannual variability in the home range of breeding turtles: implications for current and future conservation management. *Biological Conservation*, 143, 722–730.

Schofield G, Lilley MKS, Bishop CM, Brown P, Katselidis KA, Dimopoulos P, Pantis JD, Hays GC. (2010c). Conservation hotspots: Implications of intense spatial area use by breeding male and female loggerheads at the Mediterranean's largest rookery. *Endangered Species Research* 10:191–202.

Schofield G, Dimadi A, Fossette S, Katselidis KA, Koutsoubas D, Lilley MKS, Luckman A, Pantis JD, Karagouni AD, Hays GC. (2013). Satellite tracking large numbers of individuals to infer

population level dispersal and core areas for the protection of an endangered species. Diversity and Distributions 19:834– 844.

Seminoff, J.A. (Southwest Fisheries Science Center, U.S.). 2004. *Chelonia mydas*. The IUCN Red List of Threatened Species 2004: e.T4615A11037468.

Tasker, M.L., Jones, P.H., Dixon, T. & B.F. Blake (1984) Counting seabirds at sea from ships: A review of methods employed and suggestion for a standardized approach. *The Auk* 101:567-57

Ullmann, J., & M. Stachowitsch (2015) A critical review of the Mediterranean sea turtle rescue network: a web looking for a weaver.

Valeiras, J. & J. A. Caminas (2003) The incidental capture of seabirds by Spanish drifting longline fisheries in the western Mediterranean Sea. *Scientia Marina* 67: 65–68.

Velando, A. and J. Freire (1999) Inter-colony and seasonal differences in the breeding diet of European shags on the Galician coast (NW Spain). *Marine Ecology Progress Series* 188: 225–236.

Wanless, S., Harris, M.P. and J.A. Morris (1991) Foraging range and feeding locations of Shags *Phalacrocorax aristotelis* during chick rearing. *Ibis* 133: 30-36.

Temple, H.J. and Terry, A. (Compilers) (2007) *The Status and Distribution of European Mammals*. Luxembourg: Office for Official Publications of the European Communities. viii + 48pp, 210 x 297 mm.

Zbinden, J. A., Aebischer, A., Margaritoulis, D., & R. Arlettaz (2008) Important areas at sea for adult loggerhead sea turtles in the Mediterranean Sea: satellite tracking corroborates findings from potentially biased sources. *Marine Biology*, 153(5), 899-906.

Annex I: Report on the coastal surveys for the Mediterranean monk seal

Annex II: Report on Loggerhead turtles tracking in Kyparissia

Status of the important fauna species in the Kyparisiakos lease area

WP111: Coastal surveys for monk seals and Mediterranean shag breeding sites in the coastal zones of the adjacent Natura 2000 sites



Annex III. Report on the coastal surveys for the Mediterranean monk seal

November 2024

Contents

Summary.....	4
Introduction.....	5
Context of the project	5
General information on the species	5
Status and legal protection of monk seal in Greece.....	7
Importance of the project area for the Mediterranean monk seal	7
Methodology	9
Project Area	9
Collection of opportunistic monk seal observations through RINT	10
Cave Monitoring with the use of Automatic Infrared Cameras	11
Results	12
Monk seal observations collected through the operation of RINT	12
Cave Monitoring and Demographic Parameters.....	13
Data Collected through the operation of RINT	17
Conclusions.....	18
Future research	19
References.....	20

Index of Maps

Map 1 Project Area.....	9
Map 2 Caves identified during the coastal surveys of summer 2023	10
Map 3 Geographical distribution of monk seal observations collected though RINT.....	13
Map 4. The confirmed important pupping area on Zakynthos Island.....	14

Index of Photos

Photo 1: An adult male monk seal resting on a beach on the west coast of Zakynthos Island	8
Photo 2: Installation of infrared camera systems in a monk seal shelter in the project area	11
Photo 3: The earliest-born pup in cave ZAK6 suckling from its mother along two other adult females and a sub-adult.....	15
Photo 4: Three more pups born in ZAK6 visible along animals of other age classes	15
Photo 5: An adult male resting on an open beach at the area of Navagio (4-5-2024)	16
Photo 6: Three adult females with visible characteristic scars in ZAK6	16
Photo 7: Animals of all age classes photographed in ZAK6.....	17
Photo 8: A sub adult seal resting at Porto Romano (16-3-2024)	17
Photo 8: Mother and pup captured in ZAK7	18
Photo 9: Adult male captured in ZAK7 with identifiable white patch visible.....	18

Project team

Name	Expertise	Role
Panagiotis Dendrinios	PhD Biologist, Monk Seal Expert	Field Research Coordinator
Styliani Adamantopoulou	Biologist, Monk Seal Expert	Data Analysis
Kimon Koemtzopoulos	MSc. Biologist, Marine Mammal Specialist	Field Research & Data Compilation
Panagiotis Mpatzios	System Engineer/Field Technician	Field Research and technical coordination
Odysseas Paxinos	Msc, Oceanographer, Seal Specialist	Field Research & Data Compilation

Summary

The Mediterranean monk seal (*Monachus monachus*) is the rarest extant member of the Phocidae family and one of the rarest marine mammals in the world. It has been recently re-classified as “Vulnerable”, down from “Endangered” by the International Union for the Conservation of Nature. Many challenges to its survival still remain. The specific aim of the present study is the monitoring of the monk seal population along the coastline of the Kyparisiakos Gulf area in the Ionian Sea, and more specifically, along the coast of Zakynthos Island situated within the sphere of influence of future drilling operations. The area has been hypothesized to be an important breeding hub for the Mediterranean monk seal, based on preliminary data.

During the previous stages of the project, the coastline within the Wider Project Area was surveyed for the presence of suitable monk seal pupping habitat (marine caves). Monitoring efforts of the monk seal population during the previous reporting period focused on the island of Zakynthos. Three automatic infrared camera systems were placed in 2 pupping caves selected in the island of Zakynthos: ZAK6 and ZAK7. As Zakynthos island’s caves and the monk seal population using them are the most vulnerable in relation to any future drilling activities and potential spills in the Kyparisiakos lease area, monitoring of these caves was considered essential.

The present report focuses on the results of the infrared cameras placed in the aforementioned caves in combination with data collected through the operation of the Rescue and Information Network (RINT) of MOm in the area.

The results of this effort to date are summarized up as follows:

- **21 monk seal observations collected through RINT**
- **261 days of cave camera monitoring conducted**
- **13329 photographs collected**
- **17 different individuals identified through the analysis of the photos captured**
 - **7 Adult females**
 - **2 Adult males**
 - **2 Sub-adults**
 - **1 Female pup**
 - **3 Male pups**
 - **2 pups of unknown sex**

Based on the above-mentioned information, **the number of monk seals** inhabiting the area of Zakynthos Island is estimated to be **at least 20 individuals**

These results **documented for the first time the importance of Zakynthos Island as a monk seal pupping site in the Ionian region.**

Introduction

Context of the project

In the context of Environmental Monitoring and Recording of Critical Environmental Indicators of Biodiversity, such as marine mammals (cetaceans and monk seals), sea turtles and seabirds, the Hellenic Petroleum Exploration and Production of Hydrocarbons Ionian Single Member Societe Anonyme (HELPE IONIAN S.A.) company has assigned to the company Nature Conservation Consultants (NCC) Ltd, a contract for conducting the present **Project**, namely the “Survey of the Status of Important fauna species in the Kyparisiakos lease Area”.

In the framework of the Project, the project team, consisting of experienced biodiversity experts from the NCC, MOM and BIOTOPIA, implemented field surveys, at the Kyparisiakos lease Area (marine and coastal areas), which is defined as the Project Area (PA), using appropriate field sampling techniques, in order to provide sufficient and documented data on the status of marine mammals, seabirds and sea turtles, as well as other sensitive biodiversity elements and locations that should be prioritized for a future biodiversity monitoring program.

This review of the marine environment is expected to contribute to updating the existing data for the Mediterranean Monk Seal, cetaceans, seabirds and sea turtle populations, and other protected and / or threatened species in this part of the Ionian Sea.

The present document consists of the latest results of the monitoring effort for the Mediterranean monk seal using Infrared Cameras placed in previously identified (potential) pupping caves for the species on the island of Zakynthos, considered to be the main breeding hub of the species in the wider Kyparisiakos Gulf region and one of the main breeding hubs in the Ionian Sea.

General information on the species

The Mediterranean monk seal (*Monachus monachus*) is the rarest extant member of the Phocidae family and one of the rarest marine mammals in the world. It has been recently re-classified as “Vulnerable” (down from “Endangered”) by the International Union for the Conservation of Nature (Karamanlidis & Dendrinos, 2015) and is strictly protected by the Council of European Communities Directive 92/43/EEC, the Bonn, Bern, CITES, Barcelona and Rio Conventions, as well as by the Greek law (Karamanlidis et al., 2016a).

Mediterranean monk seals were once widely and continuously distributed in the Mediterranean and Black Seas, and in the North Atlantic waters from Morocco to Cap Blanc, including the Canary, Madeira and the Azores Islands. A few individuals have been recorded in Senegal, the Gambia and the Cape Verde Islands in

the southern end, as well as in Portugal and Atlantic France in the northern end of the species' distribution. Today the distribution of the Mediterranean is highly fragmented and consists of three to four isolated subpopulations (Karamanlidis et al., 2016a). In the Mediterranean Sea, the stronghold of the species has been on islands in the Ionian and Aegean Seas, and along the coasts of Greece and western and southern Turkey (Güçlüsoy, Kiraç, Ververi, & Savaş, 2004; Gücü, Gücü, & Örek, 2004; Anonymous, 2007). In the North Atlantic, two subpopulations exist: one at Cabo Blanco (also known as Cap Blanc) at the border of Mauritania and Western Sahara (González & Fernandez de Larrinoa, 2012; Martínez-Jauregui et al., 2012), and one at the Archipelago of Madeira (Pires, Neves, & Karamanlidis, 2008). An unknown number of monk seals might still survive at the Mediterranean coasts of eastern Morocco (and perhaps Algeria) (Mo, Bazairi, Bayed, & Agnesi, 2011), but without on-going systematic conservation actions the fate of this subpopulation is unknown.

A number of threats jeopardize the survival of the Mediterranean monk seal (Karamanlidis et al., 2016a). The most important are:

- **Habitat deterioration and loss** by coastal development, including disturbance by tourism and leisure boating.
- **Accidental entanglement** in fishing gear leading to death by drowning.
- **Deliberate killing** by fishermen, who consider the animal a pest that damages their nets and 'steals' their fish.
- **Other stochastic events**, such as disease outbreaks.

The Mediterranean monk seal is particularly sensitive to human disturbance, with coastal development and tourism pressures driving the species to inhabit increasingly marginal and unsuitable habitat. In some marine caves that the females use for pupping, pups are vulnerable to storm surges and may be washed away and drowned.

Conservation of the Mediterranean monk seal has been underway since the late 1970s (Johnson & Lavigne, 1998) and has focused mainly on the *in-situ* protection of the species. *In situ* conservation efforts include the establishment of marine protected areas, rescue and rehabilitation of orphaned and wounded seals, environmental education and public awareness. By providing additional knowledge into little understood aspects of the monk seal's biology and behavior scientific research (Formigaro et al., 2016; Karamanlidis et al., 2016b; Charrier, Marchesseau, Dendrinou, Tounta, & Karamanlidis, 2017), plays also a key role in promoting *in situ* conservation efforts.

There is a consensus of scientific opinion that a network of protected and well-managed sites is essential for the survival of the species (Adamantopoulou et al., 2000). To date, marine protected areas for the species have been established in only a fraction of areas throughout the current species distribution.

Status and legal protection of monk seal in Greece

Greece has been traditionally one of the species strongholds in the eastern Mediterranean and a focal point of monk seal research and conservation during the last fifty years. The first studies on the species date back to the 1970's (Marchessaux & Duguy, 1977; Marchessaux, 1979; Vamvakas, Tsimenidis, & Kainadas, 1979) and intensified over the next two (Verriopoulos, 1985; Vlachoutsikou & Lazaridis, 1990; Panou, Jacobs, & Panos, 1993; Adamantopoulou, Androukaki, & Kotomatas, 1999; Panou et al, 1999), documenting the status of the species and informing conservation actions. Recently, encouraging signs of population recovery being observed. Legal protections afforded to the species include: Mediterranean monk seal *Monachus monachus* is listed in the Red Data Book of Greece and is strictly protected by the Presidential Decree 67/1981. Organized actions to protect the species began in the country in the late 1980s. These included the establishment of protected areas, the organization of a National Rescue and Information Network and a seal care and rehabilitation program, as well as systematic environmental education in schools. The creation of the National Marine Park of Alonissos, Northern Sporades in 1992, the first Marine Protected Area established in Greece, was a milestone in monk seal conservation history in the country. Following the identification of additional important areas for the species in the country, most important pupping areas for the species have been already included in the Natura 2000 network. Today, the Mediterranean Monk Seal, which was previously categorized as “Critically Endangered” in the IUCN Red List (Aguilar & Lowry, 2008), has been showing significant signs of population recovery and was re-assessed as “Endangered” in 2015 (Karamanlidis & Dendrinis, 2015) and, following the most recent data including evidence of a significant range increase in Greece (Adamantopoulou et al, 2022), the Mediterranean Monk Seal was re-assessed as “Vulnerable” in 2023 (Karamanlidis et al, 2023). As part of the wider conservation goals for the species in Greece, the Hellenic State officially adopted the National Action Plan in 2024 for the monk seal as proposed by MOm (Dendrinis et al, 2020). The action plan includes various actions to further promote the protection of the species.

Importance of the project area for the Mediterranean monk seal


While the Kyparisiakos Gulf area is not considered as a particularly important area for the Mediterranean monk seal in the Ionian Sea, the island of Zakynthos (and especially the W and SW coasts), directly north of the study area and within the wider project area, was considered since 1990's as an important for the species site. This was, to date, based on observation reports collected through the Rescue and Information Network and on past opportunistic field visits to the island. The present research, as described below, clearly showed that along the SW coasts of Zakynthos there are very important monk seal pupping sites.

Photo 1.: An adult male monk seal resting on a beach on the west coast of Zakynthos Island

Methodology

Project Area

The **Project Area** is located in the Ionian Sea, southeast of Zakynthos Island and west of Peloponnese, More specifically it extends between latitudes of 36°50'N in the south and 37°30'N in the north and between longitudes of 20°55'E in the west and 21°30'E in the east. Its total surface area is 3,422.5 km² (Figure 1)



Map 1 Project Area

The **Wider Project Area** envelops the project area and extends further north and east to additionally include the south-western, south-eastern and eastern coast of Zakynthos, and the western coast of Peloponnese south of Kyllini, together with their neighbouring islets (Figure 1). The **Pelagic Survey Area**, where pelagic surveys are carried out includes the primarily the Project Area together with neighbouring waters to the east.

Zakynthos Island

The part of the project presented in this report, focused on the island of Zakynthos and the potential monk seal pupping caves identified during the coastal surveys conducted during 2023 (see caves distribution on Map 2).



Map 2 Caves identified during the coastal surveys of summer 2023

Collection of opportunistic monk seal observations through RINT

During the whole project period RINT was continuous operating in order to collect any available monk seal observation conducted opportunistic by citizens throughout wider project area. MOM maintains a 24/7 telephone hotline in order to collect information related to sightings of individuals of the species by citizens. At the same time, it has a special electronic form on its website (www.mom.gr) for entering data related to monk seal sightings

Cave Monitoring with the use of Automatic Infrared Cameras

During late summer 2023, 4 Infrared Camera Traps were placed in 2 selected caves at the SW coast of Zakynthos: ZAK6 and ZAK7. Three camera systems were installed in cave ZAK6 and 1 was installed in cave ZAK7. The infrared cameras were secured on the rocky walls of the caves and were positioned in a way to maximize the coverage of the terrestrial component of the cave (internal beach). Three of the camera systems were set to operate on a time lapse mode (taking pictures every 2 hours) of the cave beach to record the presence or absence of animals while one camera system set on motion detection operation mode. During Spring 2024 the infrared camera systems were retrieved and all data collected were analyzed by MOm experts.



Photo 2: Installation of infrared camera systems in a monk seal shelter in the project area

Results


Monk seal observations collected through the operation of RINT

From the 1st of January until the end of October 2024, **21 monk seal opportunistic monk seal observations** were collected through the operation of RINT. The observation details are presented in the following Table, while their geographical distribution is presented in Map 2.

Table 1 Monk seal observation collected through RINT

	Date	Area	Stage	Sex	Behavior
1	3/16/2024	Zakynthos	WEANER	Unknown	Resting
2	19/5/2024	Zakynthos	UNKNOWN	Unknown	Swimming
3	19/5/2024	Zakynthos	ADULT	Male	Resting
4	19/5/2024	Kyparissiakos Gulf	UNKNOWN	Unknown	Swimming
5	19/5/2024	Kyparissiakos Gulf	UNKNOWN	Unknown	Swimming
6	19/5/2024	Kyparissiakos Gulf	UNKNOWN	Unknown	Swimming
7	23/5/2024	Kyparissiakos Gulf	UNKNOWN	Unknown	Swimming
8	12/8/2024	Zakynthos	UNKNOWN	Unknown	Swimming
9	18/8/2024	Zakynthos	ADULT	Male	Swimming
10	19/8/2024	Sapientza	UNKNOWN	Unknown	Swimming
11	19/8/2024	Zakynthos	SUBADULT	Unknown	Resting
12	5/9/2024	Zakynthos	ADULT	Female	Swimming
13	5/9/2024	Zakynthos	ADULT	Female	Swimming
14	5/9/2024	Zakynthos	ADULT	Female	Resting
15	5/9/2024	Zakynthos	ADULT	Female	Resting
16	5/9/2024	Zakynthos	PUP1	Unknown	Sleeping
17	5/9/2024	Zakynthos	PUP1	Unknown	Sleeping
18	5/9/2024	Zakynthos	PUP1	Unknown	Resting
19	27/9/2024	Zakynthos	ADULT	Female	Swimming
20	27/9/2024	Zakynthos	ADULT	Female	Swimming
21	27/9/2024	Zakynthos	PUP1	Unknown	Resting

Extremely important are the observations regarding the presence of two newborns (Observations No17 and 18 on the 5th of September 2024) in the same cave where reproduction was also recorded in 2023 through the use of infrared cameras. These observations strongly suggested that **the reproductive activity in this particular geographical location (see Map 3) is continuous over the years and with a significant number of newborns.**



Map 3 Geographical distribution of monk seal observations collected through RINT

Cave Monitoring and Demographic Parameters

The cameras were collected during the current project period (in April 2024) and the photographs were analyzed for the presence of seals and the identification of individuals. The number of pups born in the monitored caves were also counted. The cameras that were set on a time lapse recorded images continuously from 29/7/2023 to 14/4/2024 for a total of 261 days. The second camera installed in ZAK6 and set on a motion sensor mode, functioned until the 28/1/2024 when ran out of battery.

A total of **13329 photographs** were obtained. The analysis of the photographs revealed **a regular presence of seals in the caves** as well as **the birth of 4 pups, (1 female and 3 males) during the of 2023-2024 pupping**

season. All of these were **born in cave ZAK6**. The analysis also allowed for the identification of individual mature seals and the counting of the total number of individuals using these caves.



Map 4. The confirmed important pupping area on Zakynthos Island



Photo 3: The earliest-born pup in cave ZAK6 suckling from its mother along two other adult females and a sub-adult



Photo 4: Three more pups born in ZAK6 visible along animals of other age classes

A total of 2 adult males and 7 adult females were identified. In addition, at least **2 sub-adult animals** were also found to use these caves.



Photo 5: An adult male resting on an open beach at the area of Navagio (4-5-2024)



Photo 6: Three adult females with visible characteristic scars in ZAK6

Data Collected through the operation of RINT

It is worth noting that this is the first time that the importance of this area for the Mediterranean seal has been documented using infrared cameras in marine caves.

Based on the above-mentioned important findings, a first estimate of the population status in the area indicates a total number of **at least 25 animals** that use this area with an annual pup production of at least **4 pups per year**.



Photo 7: Animals of all age classes photographed in ZAK6



Photo 8: A sub adult seal resting at Porto Romano (16-3-2024)



Photo 9: Mother and pup captured in ZAK7



Photo 10: Adult male captured in ZAK7 with identifiable white patch visible

Conclusions

The results of the current Monitoring Survey for the Monk Seal population of Zakynthos Island provide an updated (and for the first time well documented) assessment of the status of this rare species in the area.

The evidence collected during the project verify Zakynthos Island as indeed being a highly important hub for the Mediterranean monk seal not only in the Kyparisiakos Gulf wider area but in the Ionian Sea in general. Specifically cave ZAK6 shows an intense use by monk seals to give birth and raise their pups.

Some of the animals, including an adult male, a sub-adult, two adult females and at least two mother-pup pairs were also captured in cave ZAK7. The image quality from this cave was not good enough to confidently identify individual seals. Cave ZAK7 seems to be used less frequently, possibly to shelter from adverse weather conditions affecting ZAK6 as seen on the cameras. Even though the pups weren't born in ZAK7, this cave still constitutes an important component of what makes this area attractive to monk seals as a pupping location.

The up-to-date findings confirm the necessity to establish prevention and response measures and a contingency plan for the protection of the pupping habitat of the monk seals on the island in the event of an oil spill accident during future drilling operations in the region.

Future research

Considering

- the fact that Mediterranean monk seal is a priority species of the EU Habitat Directive
- the importance of the area of Zakynthos and especially the SW part of the coastline of the island the caves the already confirmed importance of the cave complex ZAK6 and ZAK7
- the indications that it is very possible that more caves in this area are used by the seals for pupping
- the fact that the area is located within the wider project area

it is suggested the continuation of the monitoring of these caves using infrared camera systems and the expansion of the infrared camera network in order to get a more complete picture of the status of the monk seal population in this area. This will allow for a more robust evaluation of the possible effects of human activities (including oil drilling operations) on the local species population.

References

- Adam, P.J. (2004). *Monachus tropicalis*. *Mammalian Species*, 747, 1-9.
- Adamantopoulou, S., Karamanlidis, A. A., Dendrinis, D. and Olivier, J. 2022. Citizen science indicates significant range recovery and defines new conservation priorities for Earth's most endangered Pinniped in Greece. *Animal Conservation* 26(1): 115-125. <https://doi.org/10.1111/acv.12806>
- Adamantopoulou, S., Anagnostopoulou, K., Dendrinis, P., Kotomatas, S., Labadariou, N., Pipinis, V., Zavras, V. (2000). Conserving critical habitats for the Mediterranean monk seal in Greece through the creation of a network of protected areas. *European Research on Cetaceans*, 14, 33.
- Adamantopoulou, S., Androukaki, E. & Kotomatas, S. (1999). The distribution of the Mediterranean monk seal in Greece based on an information network. *Contributions to the Zoogeography and Ecology of the Eastern Mediterranean Region*, 1, 399-404.
- Aguilar, A. & Lowry, L. (IUCN SSC Pinniped Specialist Group). 2008. *Monachus monachus*. The IUCN Red List of Threatened Species 2008: e.T13653A4304960. Accessed on 06 November 2024.
- Anonymous (2007). Status of the population of the Mediterranean monk seal (*Monachus monachus*) in Greece. pp. 1-42. MOM/Hellenic Society for the Study and Protection of the Monk seal, Athens, Greece.
- Charrier, I., Marchesseau, S., Dendrinis, P., Tounta, E. & Karamanlidis, A.A. (2017). Individual signatures in the vocal repertoire of the endangered Mediterranean monk seal: new perspectives for population monitoring. *Endangered species research*, 32, 459-470.
- Childerhouse, S. & Gales, N. (1998). Historical and modern distribution and abundance of the New Zealand sea lion *Phocartos hookeri*. *New Zealand Journal of Zoology*, 25, 1-16.
- Dendrinis P., Karamanlidis A.A., Adamantopoulou S., Koemtzipoulos K., Komninou A., Tounta E. (2020). MOM /Hellenic Society for the Study and Protection of the Monk seal. LIFE-IP 4 NATURA: Integrated actions for the conservation and management of Natura 2000 sites, species, habitats and ecosystems in Greece. Deliverable Action A.1: Action Plan for the Mediterranean monk seal (*Monachus monachus*). Hellenic Ministry of Environment and Energy, p. 105 ; Annexes.12p
- Dendrinis, P. (2011) Contribution to the study of the Mediterranean monk seal's (*Monachus monachus*) ecology and biology at the island complex of Northern Sporades, Greece. PhD thesis, National and Kapodistrian University of Athens.
- Dendrinis, P., Karamanlidis, A.A., Kotomatas, S., Legakis, A., Tounta, E. & Matthiopoulos, J. (2007). Pupping habitat use in the Mediterranean monk seal: a long-term study. *Marine Mammal Science*, 23, 615-628.
- Dendrinis, P., Kotomatas, S. & Tounta, E. (1999). Monk seal pup production in the National Marine Park of Alonissos-N.Sporades. *Contributions to the Zoogeography and Ecology of the Eastern Mediterranean Region*, 1, 413-419.
- Dendrinis, P., Tounta, E., Kotomatas, S. & Kottas, A. (1994). Recent data on the Mediterranean monk seal population of the Northern Sporades. *Bios (Macedonia/Greece)*, 2, 11-16.
- Formigaro, C., Karamanlidis, A.A., Dendrinis, P., Marsili, L., Silvi, M. & Zaccaroni, A. (2016). Trace element concentrations in the Mediterranean monk seal (*Monachus monachus*) in the Eastern Mediterranean Sea. *Science of the Total Environment*, 576, 528-537.
- Gerrodette, T. & Gilmartin, W.G. (1990). Demographic consequences of changed pupping and hauling sites of the Hawaiian monk seal. *Conservation Biology*, 4, 423-430.
- González, L.M. & Fernandez de Larrinoa, P. (2012). Mediterranean monk seal *Monachus monachus* distribution and fisheries interactions in the Atlantic Sahara during the second half of the 20th century. *Mammalia*, 77, 41-49.
- Güçlüsoy, H., Kiraç, C.O., Ververi, N.O. & Savaş, Y. (2004). Status of the Mediterranean monk seal, *Monachus monachus* (Hermann, 1779) in the coastal waters of Turkey. *E.U. Journal of Fisheries & Aquatic Sciences*, 21, 201-210.
- Güçlüsoy, H. & Savaş, Y. (2003). Status of the Mediterranean monk seal (*Monachus monachus*) in the Foça Pilot Monk seal Conservation area, Turkey. *Zoology in the Middle East*, 28, 5-16.

- Gücü, A.C., Gücü, G. & Orek, H. (2004). Habitat use and preliminary demographic evaluation of the critically endangered Mediterranean monk seal (*Monachus monachus*) in the Cilician Basin (Eastern Mediterranean). *Biological Conservation*, 116, 417-431.
- Johnson, W.M. & Lavigne, D.M. (1998). *The Mediterranean monk seal. Conservation Guidelines. Multilingual edition*. International Marine Mammal Association Inc., Guelph, Canada.
- Karamanlidis, A.A., Androukaki, E., Adamantopoulou, S., Chatzistryrou, A., Johnson, W.M., Kotomatas, S., Dendrinis, P. (2008). Assessing accidental entanglement as a threat to the Mediterranean monk seal *Monachus monachus*. *Endangered species research*, 5, 205-213.
- Karamanlidis, A.A. & Dendrinis, P. (2015). *Monachus monachus*. *The IUCN Red List of Threatened Species 2015: e.T13653A45227543*.
- Karamanlidis, A.A., Dendrinis, P., Fernández de Larrinoa, P., Gücü, A.C., Johnson, W.M., Kiraç, C.O. & Pires, R. (2016a). The Mediterranean monk seal *Monachus monachus*: status, biology, threats, and conservation priorities. *Mammal Review*, 46, 92-105.
- Karamanlidis, A.A., Gaughran, S., Aguilar, A., Dendrinis, P., Huber, D., Pires, R., . . . Amato, G. (2016b). Shaping species conservation strategies using mtDNA analysis: The case of the elusive Mediterranean monk seal (*Monachus monachus*). *Biological Conservation*, 193, 71-79.
- Karamanlidis, A.A., Paravas, V., Trillmich, F. & Dendrinis, P. (2010). First observations of parturition and postpartum behavior in the Mediterranean monk seal (*Monachus monachus*) in the Eastern Mediterranean. *Aquatic Mammals*, 36, 27-32.
- Karamanlidis, A.A., Pires, R., Silva, N.C. & Neves, H.C. (2004). The availability of resting and pupping habitat for the critically endangered Mediterranean monk seal *Monachus monachus* in the archipelago of Madeira. *Oryx*, 38, 180-185.
- Marchessaux, D. (1979). Nouvelles donnees sur le Phoque moine de Mediterranee *Monachus monachus*, dans le Golfe de Corinthe, Grece. *Rapport Commission internationale Mer Méditerranée*, 25/26, 143-145.
- Marchessaux, D. & Duguy, R. (1977). Le phoque moine, *Monachus monachus* (Hermann, 1779), en Grèce. *Mammalia*, 41, 419-439.
- Martínez-Jauregui, M., Tavecchia, G., Cedenilla, M.A., Coulson, T., Fernández de Larrinoa, P., Muñoz, M. & González, L.M. (2012). Population resilience of the Mediterranean monk seal *Monachus monachus* at Cabo Blanco peninsula. *Marine Ecology Progress Series*, 461, 273-281.
- Mathews, E.A. & Pendleton, G.W. (2006). Declines in harbor seal (*Phoca vitulina*) numbers in Glacier Bay National Park, Alaska, 1992-2002. *Marine Mammal Science*, 22, 167-189.
- Mattlin, R.H. (1978). Pup mortality of the New Zealand fur seal. *New Zealand Journal of Ecology*, 1, 138-144.
- Mo, G., Bazairi, H., Bayed, A. & Agnesi, S. (2011). Survey on Mediterranean monk seal (*Monachus monachus*) sightings in Mediterranean Morocco. *Aquatic Mammals*, 37, 248-255.
- MOm (2005). Status report of the Mediterranean monk seal populations in Kimolos - Polyaigos and Karpathos - Saria areas., pp. 1-34. MOm/Hellenic Society for the Study and Protection of the Monk Seal, Athens.
- MOm (2008). Final Report on the Monitoring of the Status of the Population of the Monk Seal, in Karpathos and Saria. *Final report of the project "Monitoring of the protected area Karpathos - Saria" (In Greek)*, pp. 1-77. MOm/Hellenic Society for the Study and Protection of the Monk seal, Athens, Greece.
- MOm (2017). Final Report of the Monitoring of the Mediterranean monk seal population in the National Marine Park of Zakynthos. pp. 1-42. MOm/Hellenic Society for the Study and Protection of the Monk seal, Athens.
- Orsini, J.-P., Shaughnessy, P.D. & Newsome, D. (2006). Impacts of human visitors on Australian sea lions (*Neophoca cinerea*) at Carnac Island, Western Australia: Implications for tourism management. *Tourism in Marine Environments*, 3, 101-115.
- Panou, A., Alimantiri, L., Aravantinos, P. & Verriopoulos, G. (1999). Distribution of the Mediterranean Monk Seal (*Monachus monachus*) in Greece: Results of a Pan- Hellenic Questionnaire Action, 1982-1991. *Contributions to the Zoogeography and Ecology of the Eastern Mediterranean Region*, 1, 421-428.

- Panou, A., Anestis, A., Ioannou, G., Karavellas, D.P., Liontou, E., Potamitis, A., . . . Vlachoutsikou, A. (2002a). Zakynthos Island, Ionian Sea, Greece: an important habitat for the Mediterranean monk seal, *Monachus monachus*. *9th International Congress on the Zoogeography and ecology of Greece and adjacent regions*, pp. 119. The Hellenic Zoological Society, Thessaloniki, Greece.
- Panou, A., Aravantinos, P., Kokkolis, T., Kourkoulakos, S., Lekatsa, T., Minetou, L., . . . Sclavos, E. (2002b). The Mediterranean monk seal, *Monachus monachus*, in the Central Ionian Sea, Greece: Results of a long-term study. *9th International Congress on the Zoogeography and ecology of Greece and adjacent regions*, pp. 118. The Hellenic Zoological Society, Thessaloniki, Greece.
- Panou, A., Jacobs, J. & Panos, D. (1993). The endangered Mediterranean monk seal *Monachus monachus* in the Ionian Sea, Greece. *Biological Conservation*, 64, 129-140.
- Pires, R., Neves, H.C. & Karamanlidis, A.A. (2008). The Critically Endangered Mediterranean monk seal *Monachus monachus* in the archipelago of Madeira: priorities for conservation. *Oryx*, 42, 278-285.
- Samaranch, R. & González, L.M. (2000). Changes in morphology with age in Mediterranean monk seals (*Monachus monachus*). *Marine Mammal Science*, 16, 141-157.
- Stevens, M.A. & Boness, D.J. (2003). Influences of habitat features and human disturbance on use of breeding sites by a declining population of southern fur seals (*Arctocephalus australis*). *Journal of Zoology*, 260, 145-152.
- Vamvakas, C., Tsimenidis, N. & Kainadas, H. (1979). Contribution to the Knowledge of the Distribution Pattern of the Monk Seal (*Monachus monachus*) in the Greek Seas. Conservation Plan by the Establishment of Marine Parks. *First International Conference on the Mediterranean Monk Seal* (eds K. Ronald & R. Duguy), pp. 147-150. Pergamon Press, Oxford, UK, Rhodes, Greece.
- Verriopoulos, G. (1985). Comparaison entre diverses donees sur l'evolution des populations du phoque moine en Grece. *Rapp. Comm. int. Mer Medit.*, 29, 167-168.
- Vlachoutsikou, A. & Lazaridis, Y. (1990). Monk Seals in Greece. pp. 1-43 + Appendix. Radolfzell, Germany.
- Δενδρινός, Π., Καραμανλίδης, Α. & Παράβας, Α. (2009). *Monachus monachus* (Hermann, 1779). *To Κόκκινο Βιβλίο των Απειλούμενων Ζώων της Ελλάδας* (eds Α. Λεγάκις & Π. Μαραγκού), pp. 366-369. Ελληνική Ζωολογική Εταιρεία, Αθήνα.

Tracking male turtles breeding in Kyparissia Bay: October 2024 update



Prepared by ALan Rees, PhD

Field Personnel:

Field leader:	ALan Rees
Skipper:	Kostas Papaconstantinou
Field assistant (boat):	Odysseas Paxinos
Field assistant (drone)	Odysseas Stamatakis
Field assistant (land)	Konstantinos Margaris

Cover image: View of the field site, Kyparissia Harbour, taken from a drone.

Introduction

Following successful tracking of adult female loggerhead turtles nesting in Kyparissia Bay, funded by HELPE, in 2021 we successfully deployed three transmitters on males breeding in the area in 2023. The three males received top of the range transmitters that, in addition to location, recorded and stored depth and temperature time series data with a subset of data transmitted remotely via satellite. By early spring 2024, after the turtles had reached their selected foraging locations two of the transmitters were no longer sending signals.

Male turtles often migrate to breeding sites annually, which suggested we may be able to locate the turtles in Kyparissia Bay again in 2024, and therefore retrieve the tags that contain valuable data on their behaviour from the previous 12 months.

Purpose of fieldwork

Based on the potential to retrieve the transmitters from two males we arranged a brief field visit to Kyparissia Bay. To ensure the fieldtrip would have a positive outcome even if the previously tracked turtles were not present, we arranged to deploy similar transmitters on a further two breeding males. These transmitters were supplied by ARCHELON, as part of a broader study on turtles across Greece. The fieldwork therefore had two main goals:

- 1) Retrieve transmitters from male turtles tracked from 2023 to access stored data.
- 2) Deploy transmitters on a further two male turtles

Methods

To identify if the turtles from 2023 had returned to the Kyparissia in 2024 we deployed a drone (DJI Phantom 3) to systematically search the field site for evidence of turtles with transmitters. This was undertaken throughout the field period, whilst the other work was being carried out.

We acquired the two male turtles to receive transmitters using turtle rodeo technique as used in the previous fieldwork, namely a small RIB was used to approach the turtles and then they were caught by hand, secured in a net and taken ashore to receive the transmitters. Should we have seen a male turtle with a transmitter in place, we would have used the same rodeo technique to catch it to retrieve the transmitter.



Figure 1. Left to right: The RIB and drone used to spot and catch the turtles. Capturing the turtle ready to bring ashore. Tagging and other field equipment used to record details of the captured turtle.

Results

Despite multiple hours of searching with the drone, with many turtles observed, none was recognised as having a transmitter in place. We therefore concluded that the two previously tracked turtles had not returned to Kyparissia in 2024. Our survey methods were validated as the drone observed one of the 2024 turtles in the field area after it had received its transmitter.




Figure 2. Left: Three turtles observed in close proximity to each other during drone surveys. None have a transmitter in place. Right: The first turtle to receive a transmitter in 2024 observed during the drone surveys after his release.

On the deployment side, we were able to successfully catch the planned two adult male turtles and equip them with transmitters. Both turtles' transmitters continue to transmit locations and data and they were recorded making their post-breeding migrations to widely disparate foraging locations.



Figure 3. The two adult males that were successfully equipped with satellite transmitters during the April 2024 field mission.




Figure 4. Post migration tracks of the two adult males tagged in Kyparissia Bay in May 2024.

Conclusions

The fieldwork was successful in terms of the situation present, i.e. we could not catch turtles which were not there, but we were able to catch two more to deploy additional tags. The tags were successfully deployed and continue to work as desired. This work with adult males is providing novel data on their behaviour and movements, helping fill a gap in the understanding of sea turtle biology, not only in Greece but globally.